arXiv:1412.5299v1 [math.GR] 17 Dec 2014

ADDENDA TO "FOUNDATIONS OF GARSIDE THEORY"

PATRICK DEHORNOY,
with
FRANÇOIS DIGNE, EDDY GODELLE, DAAN KRAMMER, AND JEAN MICHEL

Abstract. This text consists of additions to the book "Foundations of Garside Theory", EMS Tracts in Mathematics, vol. 22 (2015)—see introduction and table of contents in arXiv:1309.0796—namely skipped proofs and solutions to selected exercises.

Chapter I: Examples

Skipped proofs

(none)

Solution to selected exercises

(none)

Chapter II: Preliminaries

Skipped proofs

Proposition II.1.18 (**collapsing invertible elements**).— *For $\mathcal{C}$ a left-cancellative category, the following conditions are equivalent:*

(i) *The equivalence relation $=^\times$ is compatible with composition, in the sense that, if g_1g_2 is defined and $g_i'=^\times g_i$ holds for $i=1,2$, then $g_1'g_2'$ is defined and $g_1'g_2'=^\times g_1g_2$ holds;*

(ii) *The family $\mathcal{C}^\times(x,y)$ is empty for $x\neq y$ and, for all g,g' sharing the same source, $g\ {}^\times\!=^\times g'$ implies $g=^\times g'$;*

(iii) *The family $\mathcal{C}^\times(x,y)$ is empty for $x\neq y$ and we have*

$$\forall x,y\in\mathcal{O}bj(\mathcal{C})\ \forall g\in\mathcal{C}(x,y)\ \forall\epsilon\in\mathcal{C}^\times(x,x)\ \exists\epsilon'\in\mathcal{C}^\times(y,y)\ (\epsilon g=g\epsilon'). \tag{II.119}$$

When the above conditions are satisfied, the equivalence relation ${}^\times\!=^\times$ is compatible with composition, and the quotient-category $\mathcal{C}/{}^\times\!=^\times$ has no nontrivial invertible element.

Proof. Assume (i). Let $\epsilon \in \mathcal{C}^\times(x,y)$. Then $\epsilon =^\times 1_x$ and $1_y =^\times 1_y$ are satisfied, and $\epsilon\, 1_y$ is defined. By (i), $1_x\, 1_y$ must be defined as well, which is possible only for $x = y$. Let now $g \in \mathcal{C}(x,y)$ and $\epsilon \in \mathcal{C}^\times(x,x)$. Then $\epsilon =^\times 1_x$ and $g =^\times g$ are satisfied, and $\epsilon\, g$ is defined. By (i), we must have $g =^\times \epsilon g$, that is, there must exist ϵ' in $\mathcal{C}(y,y)$ satisfying $\epsilon g = g\epsilon'$. So (i) implies (iii).

Assume now (ii). Let $g \in \mathcal{C}(x,y)$ and $\epsilon \in \mathcal{C}^\times(x,x)$. Then $g \mathrel{^\times\!=^\times} \epsilon g$ holds, as we can write $\epsilon\, g = \epsilon g\, 1_y$. By (ii), we deduce $g =^\times \epsilon g$, so, as above, there must exist ϵ' in $\mathcal{C}(y,y)$ satisfying $\epsilon g = g\epsilon'$. So (ii) implies (iii).

Assume now (iii). Let $g_1 \in \mathcal{C}(x,y)$, $g_2 \in \mathcal{C}(y,z)$, and assume $g_1' =^\times g_1$ and $g_2' =^\times g_2$. By assumption, there exists ϵ_i in $\mathcal{C}^\times$ satisfying $g_i' = g_i\epsilon_i$ for $i = 1, 2$. Applying (II.1.19) to g_2 and ϵ_1, we deduce that there exists ϵ_1' in $\mathcal{C}^\times(z,z)$ satisfying $\epsilon_1 g_2 = g_2\epsilon_1'$. We deduce $g_1'g_2' = g_1g_2\epsilon_1'\epsilon_2$, whence $g_1'g_2' =^\times g_1g_2$. So (iii) implies (i).

Next, assume (iii) again, and $g' \mathrel{^\times\!=^\times} g$. By definition, there exist ϵ, ϵ' satisfying $\epsilon g' = g\epsilon'$. By (iii), there exists an invertible element ϵ'' satisfying $\epsilon g' = g'\epsilon''$, and we deduce $g' = g\,\epsilon'\epsilon''^{-1}$, whence $g' =^\times g$. So (iii) implies (ii).

Finally, assume that (i), (ii), and (iii) are satisfied, we have $g_1 \mathrel{^\times\!=^\times} g_1'$ and $g_2 \mathrel{^\times\!=^\times} g_2'$, and g_1g_2 and $g_1'g_2'$ exist. By assumption, there exist for $i = 1, 2$ invertible elements ϵ_i, ϵ_i' satisfying $\epsilon_i' g_i = g_i'\epsilon_i$. Let x and y be the source and target of g_2. By construction, $\epsilon_1^{-1}\epsilon_2'$ belongs to $\mathcal{C}^\times(x,x)$. Applying (iii), we deduce the existence of ϵ in $\mathcal{C}^\times(y,y)$ satisfying $\epsilon_1^{-1}\epsilon_2' g_2 = g_2\epsilon$. Then we obtain

$$\epsilon_1'\, g_1g_2 = \epsilon_1'\, g_1g_2\, \epsilon = \epsilon_1' g_1 \epsilon_1^{-1}\, \epsilon_2' g_2 = g_1'g_2'\, \epsilon_2,$$

which shows that $g_1g_2 \mathrel{^\times\!=^\times} g_1'g_2'$ is true. So $\mathrel{^\times\!=^\times}$ is a congruence, and there exists a well defined quotient category $\mathcal{C}/\mathrel{^\times\!=^\times}$, obtained from $\mathcal{C}$ by identifying elements that are $\mathrel{^\times\!=^\times}$-equivalent. In the current case, according to (ii), distinct objects of $\mathcal{C}$ are never connected by invertible elements, so the collapsing only involves the elements. Then, by construction, the category $\mathcal{C}/\mathrel{^\times\!=^\times}$ has no nontrivial invertible element. □

Solution to selected exercises

Exercice 4 (**atom**).— *Assume that $\mathcal{C}$ is a left-cancellative category. Show that, for $n \geqslant 1$, every element g of $\mathcal{C}$ satisfying $\mathrm{ht}(g) = n$ admits a decomposition into a product of n atoms.*

Solution. By Proposition II.2.48, there exists a decomposition $(g_1, \dots, g_n)$ of g consisting of n non-invertible entries. The assumption that g_i is non-invertible implies $\mathrm{ht}(g_i) \geqslant 1$ for every i. Hence we must have $\mathrm{ht}(g_i) = 1$ for each i, so each factor g_i is an atom. Hence g admits a decomposition as a product of n atoms.

Exercice 5 (**unique right-mcm**).— *Assume that $\mathcal{C}$ is a left-cancellative category that admits right-mcms. Assume thatf, g are elements of $\mathcal{C}$ that admit a common right-multiple and any two right-mcms of f and g are $=^\times$-equivalent. Show that every right-mcm of f and g is a right-lcm of f and g.*

Solution. Let h be a right-mcm of f and g, and $\hat{h}$ be a common right-multiple of f and g. Then $\hat{h}$ is a right-multiple of some right-mcm h' of f and g. By assumption, $h' =^\times h$ holds, hence $\hat{h}$ is a right-multiple of h as well. Hence h is a right-lcm of f and g.

Exercice 6 (**right-gcd to right-mcm**).— *Assume that $\mathcal{C}$ is a cancellative category that admits right-gcds, f, g are elements of $\mathcal{C}$ and h is a common right-multiple of f and g. Show that there exists a right-mcm h_0 of f and g such that every common right-multiple of f and g that left-divide h is a right-multiple of h_0.*

Solution. (See Figure 1.) Write $h = f\hat{g} = g\hat{f}$, and let $\hat{h}$ be a right-gcd of $\hat{f}$ and $\hat{g}$. By definition $\hat{h}$ right-divides $\hat{f}$ and $\hat{g}$, so there exist f', g' satisfying $\hat{f} = f'\hat{h}$ and $\hat{g} = g'\hat{h}$. Then we have $fg'\hat{h} = f\hat{g} = g\hat{f} = gf'\hat{h}$, whence $fg' = gf'$ by right-cancelling $\hat{h}$. Assume $fg'' = gf'' \preccurlyeq h$, say $h = fg''h''$. By left-cancelling f, we deduce $\hat{g} = g''h''$ and, similarly, $\hat{f} = f''h''$. So h'' is a common right-divisor of $\hat{f}$ and $\hat{g}$, hence it is a right-divisor of $\hat{h}$, that is, there exists h' satisfying $\hat{h} = h'h''$. This implies $fg''h'' = f\hat{g}$, whence $g''h'' = \hat{g}$ by left-cancelling f, and, finally, $h'' = fg'' \preccurlyeq f\hat{g} = h$. So every common right-multiple of f and g that left-divides h is a right-multiple of fg'.

Now assume $fg'' = gf'' \preccurlyeq fg'$. A fortiori, we have $fg'' = gf'' \preccurlyeq h$, so the above result implies $fg' \preccurlyeq fg''$, whence $fg'' =^{\times} fg'$. So fg' is a right-mcm of f and g.



FIGURE 1. Solution to Exercise 6

Exercice 8 (**conditional right-lcm**).— *Assume that $\mathcal{C}$ is a left-cancellative category.* (i) *Show that every left-gcd of fg_1 and fg_2 (if any) is of the form fg where g is a left-gcd of g_1 and g_2.* (ii) *Assume moreover that $\mathcal{C}$ admits conditional right-lcms. Show that, if g is a left-gcd of g_1 and g_2 and fg is defined, then fg is a left-gcd of fg_1 and fg_2.*

Solution. (i) Since f left-divides fg_1 and fg_2, it left-divides every left-gcd of fg_1 and fg_2, so the latter can be written fg. Now assume that fg is a left-gcd of fg_1 and fg_2. As $\mathcal{C}$ is left-cancellative, $fg \preccurlyeq fg_i$ implies $g \preccurlyeq g_i$. Next, assume that h left-divides g_1 and g_2. Then fh left-divides fg_1 and fg_2, implying $fh \preccurlyeq fg$, whence $h \preccurlyeq g$. So g is a left-gcd of g_1 and g_2.

(ii) It is clear that fg is a common left-divisor of fg_1 and fg_2. Conversely, assume that h is a common left-divisor of fg_1 and fg_2. By assumption, f and h admit fg_1 as a common right-multiple, so they admit a right-lcm, say fh'. By assumption, we have $fh' \preccurlyeq fg_1$ and $fh' \preccurlyeq fg_2$, whence $h' \preccurlyeq g_1$ and $h' \preccurlyeq g_2$ by left-cancelling f. This in turn implies $h' \preccurlyeq g$ since g is a left-gcd of g_1 and g_2. Hence we deduce $h \preccurlyeq fh' \preccurlyeq fg$, which shows that fg is a left-gcd of fg_1 and fg_2.

Exercice 9 (**left-coprime**).— *Assume that $\mathcal{C}$ is a left-cancellative category. Say that two elements f, g of $\mathcal{C}$ sharing the same source x are* left-coprime *if 1_x is a left-gcd of f and g. Assume that g_1, g_2 are elements of $\mathcal{C}$ sharing the same source, and fg_1 and fg_2 are defined. Consider the properties* (i) *The elements g_1 and g_2*

are left-coprime; (ii) *The element f is a left-gcd for fg_1 and fg_2. Show that* (ii) *implies* (i) *and that, if $\mathcal{C}$ admits conditional right-lcms,* (i) *implies* (ii)*. [Hint: Use Exercise 8.]*

Solution. If g is a non-invertible common left-divisor of g_1 and g_2, then fg is a non-invertible common left-divisor of fg_1 and fg_2, so clearly (ii) implies (i).

Conversely, assume that $\mathcal{C}$ admits conditional right-lcms and g_1, g_2 are left-coprime. By definition, 1_x is a left-gcd of g_1 and g_2. Hence, by Exercise 8, f is a left-gcd of fg_1 and fg_2. So (i) implies (ii) in this case.

Exercice 10 (**subgroupoid**).— *Let $M = \langle \mathtt{a}, \mathtt{b}, \mathtt{c}, \mathtt{d} \,|\, \mathtt{ab} = \mathtt{bc} = \mathtt{cd} = \mathtt{da} \rangle^+$ and $\Delta = \mathtt{ab}$.* (i) *Check that M is a Garside monoid with Garside element Δ.* (ii) *Let M_1 (resp. M_2) be the submonoid of M generated by* a *and* c *(resp.* b *and* c*). Show that M_1 and M_2 are free monoids of rank 2 with intersection reduced to $\{1\}$.* (iii) *Let G be the group of fractions of M. Show that the intersection of the subgroups of G generated by M_1 and M_2 is not $\{1\}$.*

Solution. (ii) No word in $\{\mathtt{a}, \mathtt{c}\}^*$ is eligible for any of the defining relations of M, so two distinct such words represent distinct elements of M_1. (iii) In G, we have $\mathtt{c}^{-1}\mathtt{a} = \mathtt{db}^{-1}$ but $\mathtt{db}^{-1}$ cannot be expressed as $f^{-1}g$ with f, g in M_2 since, otherwise, M_2 would not be free.

Exercice 11 (**weakly right-cancellative**).— *Say that a category $\mathcal{C}$ is weakly right-cancellative if $gh = h$ implies that g is invertible.* (i) *Observe that a right-cancellative category is weakly right-cancellative;* (ii) *Assume that $\mathcal{C}$ is a left-cancellative category. Show that $\mathcal{C}$ is weakly right-cancellative if and only if, for all f, g in $\mathcal{C}$, the relation $f \precsim g$ is equivalent to the conjunction of $f \precneqq g$ and "$g = g'f$ holds for no g' in $\mathcal{C}^\times$".*

Solution. (ii) The conjunction of $f \precneqq g$ and "$g = g'f$ holds for no g' in $\mathcal{C}^\times$" always implies $f \precsim g$. Assume that $\mathcal{C}$ is weakly right-cancellative and $g \succsim f$ holds. Then we have $g = g'f$ for some $g' \notin \mathcal{C}^\times$. Assume $g = g''f$: if g'' is invertible, we deduce $f = g''^{-1}g = g''^{-1}g'g$. The assumption that $\mathcal{C}$ is weakly right-cancellative implies that $g''^{-1}g'$ is invertible, hence that g' is invertible. So $f \precsim g$ implies "$g = g'f$ holds for no g' in $\mathcal{C}^\times$". Conversely, assume that $f \precsim g$ is equivalent to the conjunction of $f \precneqq g$ and "$g = g'f$ holds for no g' in $\mathcal{C}^\times$". Assume $gh = h$. If x is the source of h, we have $1_x h = h$ and 1_x is invertible. Then the assumption implies that $h \precsim h$ fails, which, as $h \preccurlyeq h$ is true, means that $h = gh$ holds for no non-invertible g.

Exercice 13 (**increasing sequences**).— *Assume that $\mathcal{C}$ is a left-cancellative. For $\mathcal{S}$ included in $\mathcal{C}$, put* $\mathrm{Div}_{\mathcal{S}}(h) = \{f \in \mathcal{C} \mid \exists g \in \mathcal{S}(fg = h)\}$*. Show that the restriction of $\precsim$ to $\mathcal{S}$ is well-founded (that is, admits no infinite descending sequence) if and only if, for every g in $\mathcal{S}$, every strictly increasing sequence in* $\mathrm{Div}_{\mathcal{S}}(g)$ *with respect to left-divisibility is finite.*

Solution. Assume that $\mathcal{S}$ is not right-Noetherian. Let $g_0, g_1, ...$ be an infinite descending sequence with respect to proper right-divisibility in $\mathcal{S}$. For each i, choose a (necessarily non-invertible) element f_i satisfying $g_{i-1} = f_i g_i$. Then we have $g_0 = f_1 g_1 = (f_1 f_2) g_2 = ...$, and the sequence 1_x (x the source of g_0), $f_1, f_1 f_2, ...$ is $\prec$-increasing in $\mathrm{Div}_{\mathcal{S}}(g_0)$.

Conversely, assume that g_0 lies in $\mathcal{S}$ and $h_1 \prec h_2 \prec ...$ is a strictly increasing sequence in $\mathrm{Div}_{\mathcal{S}}(g_0)$. Then, for each i, there exists a non-invertible element f_i

satisfying $h_i f_i = h_{i+1}$. On the other hand, as h_i belongs to $\mathrm{Div}_{\mathcal{S}}(g_0)$, there exists g_i in $\mathcal{S}$ satisfying $h_i g_i = g_0$. We find $g_0 = h_i g_i = h_{i+1} g_{i+1} = h_i f_i g_{i+1}$. By left-cancelling h_i, we deduce $g_i = f_i g_{i+1}$, hence g_{i+1} is a proper right-divisor of g_i for each i. So the sequence $g_0, g_1, ...$ witnesses that $\mathcal{S}$ is not right-Noetherian.

Exercice 15 (**left-generating**).— *Assume that $\mathcal{C}$ is a left-cancellative category that is right-Noetherian. Say that a subfamily $\mathcal{S}$ of $\mathcal{C}$* left-generates *(resp.* right-generates*) $\mathcal{C}$ if every non-invertible element of $\mathcal{C}$ admits at least one non-invertible left-divisor (resp. right-divisor) belonging to $\mathcal{S}$.* (i) *Show that $\mathcal{C}$ is right-generated by its atoms.* (ii) *Show that, if $\mathcal{S}$ is a subfamily of $\mathcal{C}$ that left-generates $\mathcal{C}$, then $\mathcal{S} \cup \mathcal{C}^\times$ generates $\mathcal{C}$.* (iii) *Conversely, show that, if $\mathcal{S} \cup \mathcal{C}^\times$ generates $\mathcal{C}$ and $\mathcal{C}^\times \mathcal{S} \subseteq \mathcal{S}^\sharp$ holds, then $\mathcal{S}$ left-generates $\mathcal{C}$.*

Solution. (ii) Assume that $\mathcal{S}$ left-generates $\mathcal{C}$. Let g be an arbitrary element of $\mathcal{C}$. If g is invertible, g belongs to $\mathcal{C}^\times$. Otherwise, by assumption, there exist a non-invertible element g_1 in $\mathcal{S}$ and g' in $\mathcal{C}$ satisfying $g = g_1 g'$. If g' is invertible, g belongs to $\mathcal{S}\mathcal{C}^\times$. Otherwise, there exist a non-invertible element g_2 in $\mathcal{S}$ and g'' satisfying $g' = g_2 g''$, and so on. By Proposition II.2.28, the sequence 1_x, g_1, $g_1 g_2$, $\cdots$, which is increasing with respect to proper left-divisibility and lies in $\mathrm{Div}(g)$, must be finite, yielding ℓ and $g = g_1 \cdots g_\ell \epsilon$ with $g_1, ..., g_\ell$ in $\mathcal{S}$ and ϵ in $\mathcal{C}^\times$.

(iii) Assume that $\mathcal{S} \cup \mathcal{C}^\times$ generates $\mathcal{C}$ and $\mathcal{C}^\times \mathcal{S} \subseteq \mathcal{S}^\sharp$ holds. Let g be a non-invertible element of $\mathcal{C}$. Let $(g_1, ..., g_p)$ be a decomposition of g such that g_i lies in $\mathcal{S} \cup \mathcal{C}^\times$ for every i. As g is not invertible, there exists i such that g_i is not invertible. Assume that i is minimal with this property. Then $g_1 \cdots g_{i-1}$ is invertible and, as $\mathcal{C}^\times \mathcal{S} \subseteq \mathcal{S}^\sharp$ holds, there exists g' in $\mathcal{S} \setminus \mathcal{C}^\times$ and ϵ in $\mathcal{C}^\times$ satisfying $g_1 \cdots g_i = g' \epsilon$. Then g' is a non-invertible element of $\mathcal{S}$ left-dividing g. Hence $\mathcal{S}$ left-generates $\mathcal{C}$.

Exercice 20 (**equivalence**).— *Assume that $(\mathcal{S}, \mathcal{R})$ is a category presentation. Say that an element s of $\mathcal{S}$ is $\mathcal{R}$-right-invertible if $sw \equiv^+_{\mathcal{R}} \varepsilon_x$ (x the source of s) holds for some w in $\mathcal{S}^*$. Show that a category presentation $(\mathcal{S}, \mathcal{R})$ is complete with respect to right-reversing if and only if, for all u, v in $\mathcal{S}^*$, the following are equivalent:* (i) *u and v are $\mathcal{R}$-equivalent (that is, $u \equiv^+_{\mathcal{R}} v$ holds),* (ii) *$\overline{u}v \curvearrowright_{\mathcal{R}} v'\overline{u'}$ holds for some $\mathcal{R}$-equivalent paths u', v' in $\mathcal{S}^*$ all of which entries are $\mathcal{R}$-right-invertible.*

Solution. Assume that right-reversing is complete for $(\mathcal{S}, \mathcal{R})$ and $u \equiv^+_{\mathcal{R}} v$ holds. Denoting by y the common target of u and v, we have $u\varepsilon_y \equiv^+_{\mathcal{R}} v\varepsilon_y$. Hence, by definition of completeness, there exist u', v', w satisfying $\overline{u}v \curvearrowright_{\mathcal{R}} v'\overline{u'}$, $\varepsilon_y \equiv^+_{\mathcal{R}} u'w$, and $\varepsilon_y \equiv^+_{\mathcal{R}} u'w$. Hence, in $\mathcal{C}$, we have $[u']^+ [w]^+ = 1_y = [v']^+ [w]^+$, whence $[u']^+ = [v']^+ = ([w]^+)^{-1}$. So u' and v' are $\mathcal{R}$-equivalent and, as their classes are invertible, they must consist of invertible entries.

Conversely, assume that the condition of (ii) is satisfied, and that uv' and vu' are $\mathcal{R}$-equivalent. By (ii), there exist $\mathcal{R}$-equivalent $\mathcal{S}$-paths u_0, v_0 such that $\overline{(uv')}(vu')$ is right-reversible to $v_0\overline{u_0}$ and all entries in u_0 and v_0 are invertible. By Lemma II.4.23, the reversing of $\overline{(uv')}(vu')$ to $v_0\overline{u_0}$ splits into four reversings. By construction, all entries in u_1, u_2, v_1, v_2 are invertible. Put $w = u'''v_2\overline{u_2 u_1}$. By assumption, we have $v_1 v_2 \equiv^+_{\mathcal{R}} u_1 u_2$, whence $\overline{v_2 v_1} \equiv^+_{\mathcal{R}} \overline{u_2 u_1}$. We deduce

$$u' \equiv^+ u''u'''v_2\overline{u_2 u_1} = u''w, \text{ and } v' \equiv^+ v''u'''v_2\overline{v_2 v_1} \equiv^+ u''w,$$

which means that (u, v, u', v') factorizing through right-reversing. Hence right-reversing is complete for $(\mathcal{S}, \mathcal{R})$.



FIGURE 2. Solution to Exercise 20

Exercice 22 (**complete** *vs.* **cube, complemented case**).— *Assume that* $(\mathcal{S},\mathcal{R})$ *is a right-complemented presentation, associated with the syntactic right-complement* θ*. Show the equivalence of the following three properties:* (i) *Right-reversing is complete for* $(\mathcal{S},\mathcal{R})$*;* (ii) *The map* θ^* *is compatible with* $\equiv^+_{\mathcal{R}}$*-equivalence in the sense that, if* $u' \equiv^+_{\mathcal{R}} u$ *and* $v' \equiv^+_{\mathcal{R}} v$ *hold, then* $\theta^*(u',v')$ *is defined if and only if* $\theta(u,v)$ *is and, if so, they are* $\equiv^+_{\mathcal{R}}$*-equivalent;* (iii) *The* θ*-cube condition is true for every triple of* $\mathcal{S}$*-paths. [Hint: For* (i) $\Rightarrow$ (ii)*, note that, if* $\theta^*(u,v)$ *is defined and* $u' \equiv^+_{\mathcal{R}} u$ *and* $v' \equiv^+_{\mathcal{R}} v$ *hold, then* $(u',v,'\theta^*(u,v),\theta^*(v,u))$ *must be* $\curvearrowright$*-factorable; for* (ii) $\Rightarrow$ (iii)*, compute* $\theta^*(w,u\theta^*(u,v))$ *and* $\theta^*(w,v\theta^*(v,u))$*; for* (iii) $\Rightarrow$ (i)*, use Lemma II.4.61.]*

Solution. (i)$\Rightarrow$(ii) Assume that $\theta^*(u,v)$ exists and $u' \equiv^+_{\mathcal{R}} u$ and $v' \equiv^+_{\mathcal{R}} v$ hold. By Corollary II.4.36, we have $u\theta^*(u,v) \equiv^+_{\mathcal{R}} v\theta^*(v,u)$, whence $u'\theta^*(u,v) \equiv^+_{\mathcal{R}} v'\theta^*(v,u)$. By assumption, the quadruple $(u',v',\theta^*(u,v),\theta^*(v,u))$ is $\curvearrowright$-factorable: this means that $\theta^*(u',v')$ and $\theta^*(v',u')$ are defined and there exists w satisfying

$$\theta^*(u,v) \equiv^+_{\mathcal{R}} \theta^*(u',v')w \quad \text{and} \quad \theta^*(v,u) \equiv^+_{\mathcal{R}} \theta^*(v',u')w.$$

By a symmetric argument, we obtain the existence of w' satisfying

$$\theta^*(u',v') \equiv^+_{\mathcal{R}} \theta^*(u,v)w' \quad \text{and} \quad \theta^*(v',u') \equiv^+_{\mathcal{R}} \theta^*(v,u)w'.$$

Merging, we deduce $\theta^*(u,v) \equiv^+_{\mathcal{R}} \theta^*(u,v)w'w$. By Corollary II.4.45, the category $\langle \mathcal{S} \mid \mathcal{R}\rangle^+$ is left-cancellative, so we deduce $\varepsilon_- \equiv^+_{\mathcal{R}} w'w$, whence $w' = w = \varepsilon_-$ and, finally, $\theta^*(v',u') \equiv^+_{\mathcal{R}} \theta^*(u,v)$.

(ii)$\Rightarrow$(iii) Assume that θ^* is compatible with $\equiv^+_{\mathcal{R}}$ and $\theta^*_3(u,v,w)$ is defined. By Lemma II.4.6, we have $\theta^*_3(u,v,w) = \theta^*(u(RC^*(u,v),w)$. As $u(\theta^*(u,v) \equiv^+_{\mathcal{R}} v\theta^*(v,u)$ holds, the compatibility assumption implies that $\theta^*(v(RC^*(v,u),w)$ is defined as well, and it is $\equiv^+_{\mathcal{R}}$-equivalent to the latter. Now, by Lemma II.4.6 again, the path $\theta^*(v(\theta^*(v,u),w)$ is $\theta^*_3(v,u,w)$, so we conclude that $\theta^*_3(v,u,w)$ is defined and it is $\equiv^+_{\mathcal{R}}$-equivalent to $\theta^*_3(u,v,w)$, that is, the θ-cube condition is true for (u,v,w).

(iii)$\Rightarrow$(i) Lemma II.4.61 gives the result directly.

Exercice 23 (**alternative proof**).— *Assume that* $(\mathcal{S},\mathcal{R})$ *is a right-complemented presentation associated with the syntactic right-complement* θ*, that* $(\mathcal{S},\mathcal{R})$ *is right-Noetherian, and that the* θ*-cube condition is true on* $\mathcal{S}$*.* (i) *Show that, for all* r,s,t *in* $\mathcal{S}$*, the path* $\theta^*(r,s\theta(s,t))$ *is defined if and only if* $\theta^*(r,t\theta(t,s))$ *is and, in this case, the relations* $\theta^*(r,s\theta(s,t)) \equiv^+_{\mathcal{R}} \theta^*(r,t\theta(t,s))$ *and* $\theta^*(s\theta(s,t),r) \equiv^+_{\mathcal{R}} \theta^*(t\theta(t,s),r)$ *are satisfied.* (ii) *Show that the map* θ^* *is compatible with* $\equiv^+_{\mathcal{R}}$*, that is, the conjunction of* $u' \equiv^+_{\mathcal{R}} u$ *and* $v' \equiv^+_{\mathcal{R}} v$ *implies that* $\theta^*(u',v')$ *exists if and only if* $\theta^*(u,v)$ *does and, in this case, they are* $\equiv^+_{\mathcal{R}}$*-equivalent. [Hint: Show using on* $\lambda^*(u\theta^*(u,v))$*, where* λ^* *is a right-Noetherianity witness for* $(\mathcal{S},\mathcal{R})$ *that, if* $\theta^*(u,v)$ *is defined and we have* $u' \equiv^+_{\mathcal{R}} u$ *and* $v' \equiv^+_{\mathcal{R}} v$*, then* $\theta^*(u',v')$ *is defined and we have* $\theta^*(u',v') \equiv^+_{\mathcal{R}} \theta^*(u,v)$

and $\theta^*(v',u') \equiv^+_{\mathcal{R}} \theta^*(v,u)$*.]* (iii) *Apply Exercise 22 to deduce a new proof of Proposition II.4.16 in the right-Noetherian case.*

Solution. (i) Lemma II.4.6 gives

$$\theta^*(s\theta(s,t),r) = \theta^*(\theta(s,t),\theta(s,r)) = \theta^*_3(s,t,r),$$
$$\theta^*(r,s\theta(s,t)) = \theta(r,s)\theta^*(\theta(s,r),\theta(s,t)) = \theta(r,s)\theta^*_3(s,r,t).$$

As the θ-cube condition is true on $\{r,s,t\}$, we have $\theta^*_3(s,t,r) \equiv^+_{\mathcal{R}} \theta^*_3(t,r,r)$, so, using the first equality above and its counterpart exchanging s and t, we find

$$\theta^*(s\theta(s,t),r) = \theta^*_3(s,t,r) \equiv^+_{\mathcal{R}} \theta^*_3(t,r,r) = \theta^*(t\theta(t,s),r).$$

Similarly, the θ-cube condition gives the relations $\theta^*_3(s,r,t) \equiv^+_{\mathcal{R}} \theta^*_3(r,s,t)$ and $\theta^*_3(r,t,s) \equiv^+_{\mathcal{R}} \theta^*_3(t,r,s)$, whence, by Corollary II.4.36,

$$\begin{aligned}\theta^*(r,s\theta(s,t)) = \theta(r,s)\theta^*_3(s,r,t) &\equiv^+_{\mathcal{R}} \theta(r,s)\theta^*_3(r,s,t)\\ &\equiv^+_{\mathcal{R}} \theta(r,t)\theta^*_3(r,t,s) \equiv^+_{\mathcal{R}} \theta(r,t)\theta^*_3(t,r,s) = \theta^*(r,t\theta(t,s)).\end{aligned}$$

(ii) The result of (i) is the compatiblity in the case $u' = u = r$ and $v = s\theta(s,t)$, $v' = t\theta(t,s)$, that is, in the basic case of equivalence. We establish the general result using induction on $\lambda^*(u\theta^*(u,v))$ and, for a given value α, on the sum d of the combinatorial distances from u to u' and from v to v'. For an induction, it is sufficient to consider the case $d = 1$, that is, we may assume $u' = u$ and $\mathrm{dist}(v,v') = 1$, that is, there exist s,t in $\mathcal{S}$ and v_0, v_1 in $\mathcal{S}^*$ satisfying $v = v_0 s\theta(s,t)v_1$ and $v' = v_0 t\theta(t,s)v_1$. We assume that $\theta^*(u,v)$ is defined, and our aim is to show that $\theta^*(u',v')$ is defined as well and we have $\theta^*(u',v') \equiv^+_{\mathcal{R}} \theta^*(u,v)$ and $\theta^*(v',u') \equiv^+_{\mathcal{R}} \theta^*(v,u)$. To this end, we compare the reversing grids below:



By assumption, the left grid exists, and we wish to show that the right grid exists as well and that the corresponding paths are pairwise $\equiv^+_{\mathcal{R}}$-equivalent. The rectangles on the left (u and v_0) coincide. Next, we find a rectangle as in (i), namely r and $s\theta(s,t)$ *vs.* r and $t\theta(t,s)$. By (i), u'_2 and v'_3 exist and we have $u'_2 \equiv^+_{\mathcal{R}} u_2$ and $v'_3 \equiv^+_{\mathcal{R}} v_3$. Then consider the median bottom rectangles (u_1 and v'_3): the point is the inequality

$$\lambda^*(\theta^*(u_1v_5)) < \lambda^*(ru_1v_5) \leqslant \lambda^*(ru_1v_5v_6) \leqslant \lambda^*(v_0ru_1v_5v_6) = \lambda^*(u,v) \leqslant \alpha.$$

As $\theta^*(u_1,v_3)$ exists and we have $v'_3 \equiv^+_{\mathcal{R}} v_3$, the induction hypothesis implies that $\theta^*(u_1,v'_3)$ exists as well and gives $u'_4 \equiv^+_{\mathcal{R}} u_4$ and $v'_5 \equiv^+_{\mathcal{R}} v_5$. The argument is the same for the two right squares.

(iii) So, if the θ-cube condition is true on $\mathcal{S}$, the map θ^* is compatible with $\equiv^+_{\mathcal{R}}$. By Exercise 22, the latter condition implies that right-reversing is complete for $(\mathcal{R},\mathcal{S})$.



FIGURE 3. Solution to Exercise 24: The cube condition viewed as a factorization property: whenever $\overline{u}w\overline{w}v$ right-reverses to $\hat{v}\overline{\hat{u}}$, the quadruple $(u, v, \hat{u}, \hat{v})$ is $\curvearrowright$-factorable.

Exercice 24 (**cube condition**).— *Assume that $(\mathcal{S}, \mathcal{R})$ is a category presentation* (i) *Show that the cube condition is true for (u, v, w) if and only if, for all $\hat{u}, \hat{v}$ in $\mathcal{S}^*$ satisfying $\overline{u}w\overline{w}v \curvearrowright_{\mathcal{R}} \hat{v}\overline{\hat{u}}$, the quadruple $(u, v, \hat{u}, \hat{v})$ is $\curvearrowright$-factorable.* (ii) *Show that, if right-reversing is complete for $(\mathcal{S}, \mathcal{R})$, then the cube condition is true for every triple of $\mathcal{S}$-paths.*

Solution. (ii) Assume that u, v, w belong to $\mathcal{S}^*$, and we have $\overline{u}w \curvearrowright_{\mathcal{R}} v_1\overline{u_0}$, $\overline{w}v \curvearrowright_{\mathcal{R}} v_0\overline{u_1}$, and $\overline{u_0}v_0 \curvearrowright_{\mathcal{R}} v_2\overline{u_2}$ (so that u, v, w necessarily share the same source). By Proposition II.4.34, we have $uv_1 \equiv^+_{\mathcal{R}} wu_0$, $vu_1 \equiv^+_{\mathcal{R}} wv_0$, and $u_0v_2 \equiv^+_{\mathcal{R}} v_0u_2$, and we deduce $uv_1v_2 \equiv^+_{\mathcal{R}} wu_0v_2 \equiv^+_{\mathcal{R}} wv_0u_2 \equiv^+_{\mathcal{R}} vu_1u_2$. The assumption that right-reversing is complete then implies that (u, v, u_1u_2, v_1v_2) is $\curvearrowright$-factorable, which exactly means that the cube condition is true for (u, v, w).

Chapter III: Normal decompositions

SKIPPED PROOFS

Proposition III.1.14 (**power**).— *If $\mathcal{S}$ is a subfamily of a left-cancellative category $\mathcal{C}$ and $g_1|\cdots|g_p$ is $\mathcal{S}$-greedy, then $g_1\cdots g_m|g_{m+1}\cdots g_p$ is $\mathcal{S}^m$-greedy for $1 \leqslant m \leqslant p$, that is,*

(III.1.15) *Each relation $s \preccurlyeq fg_1\cdots g_p$ with s in $\mathcal{S}^m$ implies $s \preccurlyeq fg_1\cdots g_m$.*

Proof. (See Figure 4.) We use induction on m. For $m = 1$, the result follows from Proposition III.1.12, and more precisely from (III.1.13). Assume $m \geqslant 2$. Let $s \in \mathcal{S}^m$, say $s = s_1\cdots s_m$ with $s_1, ..., s_m$ in $\mathcal{S}$, and $s \preccurlyeq fg_1\cdots g_p$. Then we have $s_1 \preccurlyeq fg_1\cdots g_p$, hence, by (III.1.13), $s_1 \preccurlyeq fg_1$, say $fg_1 = s_1f_1$, and, therefore, $s_2\cdots s_m \preccurlyeq f_1g_2\cdots g_p$. As $s_2\cdots s_m$ belongs to $\mathcal{S}^{m-1}$, the induction hypothesis implies $s_2\cdots s_m \preccurlyeq f_1g_2\cdots g_m$, whence $s_1s_2\cdots s_m \preccurlyeq fg_1g_2\cdots g_m$, as expected. □

Lemma III.2.52.— *For f, g, f', g' in a cancellative category $\mathcal{C}$ satisfying $f'g = g'f$, the following conditions are equivalent:*

(i) *The elements f' and g' are left-disjoint;*
(ii) *The element $f'g$ is a weak left-lcm of f and g.*



FIGURE 4. Inductive proof of Proposition III.1.14.

Proof. Assume (i), and assume that $f''g = g''f$ is a common left-multiple of f and g such that $f''g$ and $f'g$ admit a common left-multiple, say $h'(f''g) = h''(f'g)$. By assumption we also have $h'(g''f) = h''(g'f)$ and, because $\mathcal{C}$ is assumed to be right-cancellative, we deduce $h'f'' = h''f'$ and $h'g'' = h''g'$. Thus (h', h'') witnesses for $(f'', g'') \bowtie (f', g')$. As f' and g' are left-disjoint, we deduce $h' \preccurlyeq h''$, that is, there exists h satisfying $h'' = h'h$. We deduce $h'(f''g) = h''(f'g) = h'h(f'g)$, whence $f''g = h(f'g)$ by left-cancelling h'. This shows that $f''g$ is a left-multiple of $f'g$, and the latter is a weak left-lcm of f and g. So (i) implies (ii).

Assume now (ii), and assume that (h', h'') witnesses for $(f'', g'') \bowtie (f', g')$, that is, we have $h'f'' = h''f'$ and $h'g'' = h''g'$. We deduce $h'f''g = h''f'g = h''g'f = h'g''f$, whence $f''g = g''f$ by left-cancelling h'. So $f''g$ is a common left-multiple of f and g. Moreover, the above equalities show that $f''g$ and $f'g$ admit a common left-multiple. As $f'g$ is a weak left-lcm of f and g, we deduce that $f''g$ is a left-multiple of $f'g$, that is, there exists h satisfying $f''g = hf'g$, hence $h''f'g = h'f''g = h'hf'g$. Right-cancelling $f'g$, we deduce $h'' = h'h$, that is, $h' \preccurlyeq h''$. Hence f' and g' are left-disjoint, and (ii) implies (i). □

Proposition III.2.53 (**symmetric normal exist**).— *If $\mathcal{S}$ is a Garside family in a cancellative category $\mathcal{C}$, the following conditions are equivalent:*

(i) *For all f, g in $\mathcal{C}$ admitting a common right-multiple, there exists a symmetric $\mathcal{S}$-normal path $\overline{u}v$ satisfying $(f, g) \bowtie ([u]^+, [v]^+)$;*

(ii) *The category $\mathcal{C}$ admits conditional weak left-lcms.*

Proof. Assume (i), and let f, g be two elements of $\mathcal{C}$ that admit a common left-multiple, say $\hat{f}g = \hat{g}f$. By (i) applied to $\hat{f}$ and $\hat{g}$, there exists a symmetric $\mathcal{S}$-normal path $\overline{u}v$ such that, putting $f' = [u]^+$ and $g' = [v]^+$, we have $(\hat{f}, \hat{g}) \bowtie (f', g')$, that is, there exist $h', \hat{h}$ satisfying $h'\hat{f} = \hat{h}f'$ and $h'\hat{g} = \hat{h}g'$. By Proposition III.2.11, f' and g' are left-disjoint, so, by Lemma III.2.52, $f'g$, which is also $g'f$, is a weak left-lcm of f and g. Hence $\mathcal{C}$ admits conditional weak left-lcms, and (i) implies (ii).

Conversely, assume (ii), and let f, g be two elements of $\mathcal{C}$ that admit a common right-multiple, say $fg' = gf'$. By (ii), there exists a weak left-lcm of f' and g', say $f''g' = g''f'$, and h satisfying $fg' = hf''g'$, hence $f = hf''$ by right-cancelling g' and, similarly, $g = hg''$. By Lemma III.2.52, the elements f'' and g'' are left-disjoint. Let u be an $\mathcal{S}$-normal decomposition of f'' and v be an $\mathcal{S}$-normal decomposition of g''. By (the trivial part of) Proposition III.2.11, the first entries of u and v are left-disjoint since f'' and g'' are, so $\overline{u}v$ is symmetric $\mathcal{S}$-normal. Finally, the pair $(h, 1_x)$ (with x the source of f and g) witnesses for $(f, g) \bowtie (f'', g'')$. So (ii) implies (i). □

Lemma III.2.55.— *A cancellative category $\mathcal{C}$ admits conditional weak left-lcms if and only if $\mathcal{C}$ is a strong Garside family in itself.*

Proof. Assume that $\mathcal{C}$ is strong and $fs = gt$ holds. Then there exist f', g', h such that f' and g' are left-disjoint and we have $f's = h't$, $f = hf'$, and $g = hg'$. By Lemma III.2.52, $f'g$ is a weak left-lcm of f and g, of which fg is a left-multiple. So $\mathcal{C}$ admits conditional weak left-lcms.

Conversely, assume that $\mathcal{C}$ admits conditional weak left-lcms, and $fs = gt$ holds. Then there exists a weak left-lcm $f's$ of s and t of which fs is a left-multiple. By Lemma III.2.52 again, f' and g' are left-disjoint, and the condition of Definition III.2.54 is satisfied. So $\mathcal{C}$ is strong. □

Proposition III.2.56 (**symmetric normal, short case III**).— *If $\mathcal{S}$ is a strong Garside family in a cancellative category $\mathcal{C}$ admitting conditional weak left-lcms, Algorithm III.2.42 running on a positive–negative $\mathcal{S}^\sharp$-path $v\overline{u}$ such that $[u]^+$ and $[v]^+$ admit a common left-multiple, say $\hat{f}[v]^+ = \hat{g}[u]^+$, returns a symmetric $\mathcal{S}$-normal path $\overline{u''}v''$ satisfying $(f, g) \bowtie ([u'']^+, [v'']^+)$ and $[u''v]^+ = [v''u]^+$; moreover there exists h satisfying $f = h[u'']^+$ and $g = h[v'']^+$.*

Once again, Proposition III.2.56 reduces to Proposition III.2.44 in the case of a left-Ore category as it then says that $\overline{u}v$ is a decomposition of $[v\overline{u}]$ in $\mathcal{E}nv(\mathcal{C})$.

Proof. The argument is the same as for Proposition III.2.44, the only difference being that, at each left-reversing step, one has to check the existence of a factorization of the initial equality $f[v] = g[u]$. The principle is explained in Figure 5: the induction hypothesis that is maintained at each step in the construction of the rectangular diagram is that, for every local North-West corner in the current diagram, there exists a factorizing arrow coming from the top-left object x. When one more tile is added, the defining property of a strong Garside family guarantees that one can add a new tile in which the left and top arrows represent left-disjoint elements and there exists a factorizing arrow coming from x. The rest of the proof is unchanged as, in particular, the third domino rule is still valid in the extended context. □

Solution to selected exercises

Exercice 28 (**invertible**).— *Assume that $\mathcal{C}$ is a left-cancellative category and $\mathcal{S}$ is included in $\mathcal{C}$. Show that, if $g_1 \cdots g_p$ belongs to $\mathcal{S}^\sharp$, then $g_1|\cdots|g_p$ being $\mathcal{S}$-greedy implies that $g_2, \dots, g_p$ are invertible.*

Solution. The element $g_1 \cdots g_p$ lies in $\mathcal{S}^\sharp$ is equal to, hence left-divides, itself. By Proposition III.1.12, $g_1|g_2 \cdots g_p$ is $\mathcal{S}$-greedy, so we deduce that $g_1 \cdots g_p$ left-divides g_1, say $g_1 = g_1 \cdots g_p g'$. Left-cancelling g_1, we deduce that $g_2 \cdots g_p g'$ is an identity-element, hence $g_2, \dots, g_p$, and g' must be invertible.

Exercice 29 (**deformation**).— *Assume that $\mathcal{C}$ is a left-cancellative category. Show that a path $g_1|\cdots|g_q$ is a $\mathcal{C}^\times$-deformation of $f_1|\cdots|f_p$ if and only if $g_1 \cdots g_i =^\times f_1 \cdots f_i$ holds for $1 \leqslant i \leqslant \max(p, q)$, the shorter path being extended by identity-elements if needed.*

Solution. Let $r = \max(p, q)$. Assume that $\epsilon_0, \dots, \epsilon_r$ are invertible elements witnessing that $g_1|\cdots|g_r$ is a $\mathcal{C}^\times$-deformation of $f_1|\cdots|f_r$. For every i, we deduce $f_1 \cdots f_i \epsilon_i = \epsilon_0 g_1 \cdots g_r$, whence $g_1 \cdots g_i =^\times f_1 \cdots f_i$ since ϵ_0 is an identity-element.



FIGURE 5. Proof of Proposition III.2.56: when the rectangular grid of Figure III.18 is constructed, there exists at each step an arrow connecting the top–left corner to each local North-West corner of the current diagram.

Conversely, assume $g_1 \cdots g_i =^{\times} f_1 \cdots f_i$ for every i, say $g_1 \cdots g_i = f_1 \cdots f_i \epsilon_i$ with ϵ_i in $\mathcal{C}_i$. Set $\epsilon_0 = 1_x$ where x is the source of f_1. Then we have $f_1 \epsilon_1 = g_1$ by construction. Assume $i \geqslant 2$. Then we obtain $f_1 \cdots f_{i-1} f_i \epsilon_i = g_1 \cdots g_{i-1} g_i = f_1 \cdots f_{i-1} \epsilon_{i-1} g_i$, whence $f_i \epsilon_i = \epsilon_{i-1} g_i$ by left-cancelling $f_1 \cdots f_{i-1}$. So $g_1 | \cdots | g_r$ is a $\mathcal{C}^{\times}$-deformation of $f_1 | \cdots | f_r$.

Exercice 33 (**left-disjoint**).— *Assume that $\mathcal{C}$ is a left-cancellative category, f and g are left-disjoint elements of $\mathcal{C}$, and f left-divides g. Show that f is invertible.*

Solution. Let x be the common source of f and g. By assumption we have $f \preccurlyeq 1_x f$ and $f \preccurlyeq 1_x g$. By definition of f and g being left-disjoint, this implies $f \preccurlyeq 1_x$, hence f must be invertible.

Exercice 34 (**normal decomposition**).— *Give a direct argument from deriving Proposition III.2.20 from Corollary III.2.50 in the case when $\mathcal{S}$ is strong.*

Solution. Let gf^{-1} be an element of $\mathcal{C}\mathcal{C}^{-1}$. Let $s_1 | \cdots | s_p$ be an $\mathcal{S}$-normal decomposition of f, and let $t_1 | \cdots | t_q$ be an $\mathcal{S}$-normal decomposition of g. We prove the existence of an $\mathcal{S}$-normal decomposition for gf^{-1} using an induction on p to construct a rectangular diagram consisting of p rows of q tiles as in Lemma III.2.31. As $t_1 | \cdots | t_q$ is $\mathcal{S}$-normal, Corollary III.2.50 inductively implies that the elements of every horizontal line of the diagram make an $\mathcal{S}$-normal path, and so do in particular the elements $t'_1 | \cdots | t'_q$ of the top line. Similarly, as $s_1 | \cdots | s_p$ is $\mathcal{S}$-normal, Corollary III.2.50 again inductively implies that the elements of every vertical line of the diagram make an $\mathcal{S}$-normal path, and so do in particular the elements $s'_1 | \cdots | s'_p$ of the left line. Finally, s'_1 and t'_1 are left-disjoint by construction. Hence $\overline{s'_p} | \cdots | \overline{s'_1} | t'_1 | \cdots | t'_q$ is an $\mathcal{S}$-normal decomposition of gf^{-1}.

Exercice 35 (**Garside base**).— (i) *Let $\mathcal{G}$ be the category whose diagram is displayed on Figure 6 left, and let $S = \{\mathtt{a}, \mathtt{b}\}$. Show that $\mathcal{G}$ is a groupoid with nine elements, $\mathcal{S}$ is a Garside base in $\mathcal{G}$, the subcategory $\mathcal{C}$ of $\mathcal{G}$ generated by $\mathcal{S}$ contains*

no nontrivial invertible element, but $\mathcal{C}$ is not an Ore category. Conclusion? (ii) *Let $\mathcal{G}$ be the category whose diagram is displayed on Figure 6 right, let $\mathcal{S} = \{\epsilon, \mathtt{a}\}$, and let $\mathcal{C}$ be the subcategory of $\mathcal{G}$ generated by $\mathcal{S}$. Show that $\mathcal{G}$ is a groupoid and every element of $\mathcal{G}$ admits a decomposition that is symmetric $\mathcal{S}$-normal in $\mathcal{C}$. Show that $\epsilon\mathtt{a}$ admits a symmetric $\mathcal{S}$-normal decomposition and no $\mathcal{S}$-normal decomposition. Is $\mathcal{S}$ a Garside family in $\mathcal{C}$? Conclusion?*



FIGURE 6. Diagrams of the categories of Exercise 35.

Solution. (i) The nine elements of $\mathcal{G}$ are 1_0, 1_1, 1_2, $\mathtt{a}$, $\mathtt{b}$, $\mathtt{a}^{-1}$, $\mathtt{b}^{-1}$, $\mathtt{a}^{-1}\mathtt{b}$, and $\mathtt{b}^{-1}\mathtt{a}$. That $\mathcal{S}$ is a Garside base in $\mathcal{G}$ follows from a direct inspection. The subcategory of $\mathcal{G}$ generated by $\mathcal{S}$ comprises five elements: 1_0, 1_1, 1_2, $\mathtt{a}$, and $\mathtt{b}$, none of which is invertible. Next, $\mathcal{C}$ is not an Ore category since the elements $\mathtt{a}$ and $\mathtt{b}$ have non common right-multiple although they share the same source. So, in Proposition III.2.25, the conclusion cannot be strengthened to claim that $\mathcal{C}$ necessarily is an Ore category (whereas the assumption of Proposition III.2.24 cannot be weakened to only assume that $\mathcal{C}$ is a left-Ore category).

(ii) The elements of $\mathcal{G}$ are 1_0, 1_1, ϵ, $\mathtt{a}$, $\epsilon\mathtt{a}$, $\mathtt{a}^{-1}$, and $\mathtt{a}^{-1}\epsilon$; those of $\mathcal{C}$ are 1_0, 1_1, ϵ, $\mathtt{a}$, $\epsilon\mathtt{a}$. Then $\epsilon\mathtt{a}$ admits the symmetric $\mathcal{S}$-normal decomposition $\overline{\epsilon}|\mathtt{a}$, but admits no $\mathcal{S}$-normal decomposition. Hence $\mathcal{S}$ is not a Garside family in $\mathcal{C}$. So, in Proposition III.2.25, we cannot simply drop the assumption about nontrivial invertible elements.

Chapter IV: Recognizing Garside families

SKIPPED PROOFS

Lemma IV.1.13.— *Assume that $\mathcal{S}$ is a subfamily of a left-cancellative category $\mathcal{C}$.*

(ii) *The family $\mathcal{S}$ is closed under right-comultiple if and only if $\mathcal{S}^\sharp$ is.*

(iii) *If $\mathcal{S}$ is closed under right-complement and $\mathcal{C}^\times\mathcal{S} \subseteq \mathcal{S}$ holds, then $\mathcal{S}^\sharp$ is closed under right-complement too.*

Proof. (ii) Assume that $\mathcal{S}^\sharp$ is closed under right-comultiple, and we have $sg = tf$ with s, t in $\mathcal{S}$. The assumption implies the existence of s', t', h satisfying $st' = ts' \in \mathcal{S}^\sharp$, $f = s'h$, and $g = t'h$. If st' is invertible, then s and t must be invertible too, so s is a common right-multiple of s and t lying in $\mathcal{S}$ of which sg is a right-multiple. Assume now that st' is not invertible. Then there exists ϵ in $\mathcal{C}^\times$ such that $st'\epsilon^{-1}$ lies in $\mathcal{S}$. Then we have $s(t'\epsilon^{-1}) = t(s'\epsilon^{-1}) \in \mathcal{S}$, $f = (s'\epsilon^{-1})(\epsilon h)$, and $g = (t'\epsilon^{-1})(\epsilon h)$. Hence $s'\epsilon^{-1}$, $t'\epsilon^{-1}$, and ϵh witness for $\mathcal{S}$ being closed under right-comultiple.

Conversely, assume that $\mathcal{S}$ is closed under right-comultiple and we have $sg = tf$ with $s, t \in \mathcal{S}^\sharp$. Assume first that s or t is invertible, say s. Put $s' = 1_y$ where y is

the target of t, $t' = s^{-1}t$, and $h = f$. Then we have $st' = ts'$, $f = s'h$, $g = t'h$, and s' and t' are invertible, hence lie in $\mathcal{S}^\sharp$, and, by assumption, so does st', which is t. So st' is a common right-multiple of s and t that lies in $\mathcal{S}^\sharp$ and of which sg is a right-multiple. Assume now that neither s nor t is invertible. Then s and t lie in $\mathcal{S}\mathcal{C}^\times$, and there exist s', t' in $\mathcal{S}$ and ϵ, ϵ' in $\mathcal{C}^\times$ satisfying $s = s'\epsilon$ and $t = t'\epsilon'$, see Figure 7. Then we have $s'(\epsilon g) = t'(\epsilon' f)$ with s', t' in $\mathcal{S}$. As $\mathcal{S}$ is closed under right-comultiple, there must exist s'', t'', and h satisfying

$$s't'' = t's'' \in \mathcal{S}, \quad \epsilon' f = s''h, \quad \text{and} \quad \epsilon g = t''h.$$

As ϵ and ϵ' are invertible, we can put $s''_1 = \epsilon'^{-1}s''$ and $t''_1 = \epsilon^{-1}t''$. Then we have $f = s''_1 h$ and $g = t''_1 h$, and $st''_1 = s't'' = t's'' = ts''_1 \in \mathcal{S} \subseteq \mathcal{S}^\sharp$. So, again, st''_1 is a common right-multiple of s and t that lies in $\mathcal{S}^\sharp$ and of which sg is a right-multiple. Hence $\mathcal{S}^\sharp$ is closed under right-comultiple.

(iii) Assume now that $\mathcal{S}$ is closed under right-complement, $\mathcal{C}^\times\mathcal{S} \subseteq \mathcal{S}$ holds, and we have $s, t \in \mathcal{S}^\sharp$ and $sg = tf$. We follow the same scheme as for (ii), and keep the same notation. Assume first that s or t is invertible, say s. Put $s'' = 1_y$ (y the target of t, $t' = s^{-1}t$, and $h = f$. Then we have $st' = ts''$, $f = s''h$, and $g = t'h$. Moreover, s'' belongs to $\mathcal{S}^\sharp$ by definition and t', which belongs to $\mathcal{C}^\times\mathcal{S}^\sharp$, hence to $\mathcal{C}^\times \cup \mathcal{C}^\times\mathcal{S}\mathcal{C}^\times$, belongs to $\mathcal{S}^\sharp$ as $\mathcal{C}^\times\mathcal{S}$ is included in $\mathcal{S}$. Assume now that neither s nor t is invertible. Then we write $s = s'\epsilon$ and $t = t'\epsilon'$ with s', t' in $\mathcal{S}$ and ϵ, ϵ' in $\mathcal{C}^\times$, see Figure 7 again. We have $s'(\epsilon g) = t'(\epsilon' f)$ so, as $\mathcal{S}$ is closed under right-complement, there exist s'', t'' in $\mathcal{S}$ and h in $\mathcal{C}$ satisfying $s't'' = t's''$, $\epsilon g = t''h$, and $\epsilon' f = s''h$. Put $s''_1 = \epsilon'^{-1}s''$ and $t''_1 = \epsilon^{-1}t''$. Then we have $st''_1 = ts''_1$, $f = s''_1 h$, and $g = t''_1 h$. Moreover, by construction, s''_1 and t''_1 belong to $\mathcal{C}^\times\mathcal{S}$, hence, by assumption, to $\mathcal{S}^\sharp$. Hence $\mathcal{S}^\sharp$ is closed under right-complement. □



FIGURE 7. Proof of Lemma IV.1.13.

Lemma IV.2.39.— *For every subfamily $\mathcal{S}$ of a left-cancellative category $\mathcal{C}$ that is right-Noetherian and admits unique conditional right-lcms, the following conditions are equivalent:*

(i) *The family $\mathcal{S}^\sharp$ is closed under right-complement (in the sense of Definition IV.1.3);*

(ii) *The family $\mathcal{S}^\sharp$ is closed under $\backslash$, that is, if s and t belong to $\mathcal{S}^\sharp$, then so does $s\backslash t$ when defined, that is, when s and t admit a common right-multiple.*

Proof. Assume that $\mathcal{S}^\sharp$ is closed under right-complement. Let s, t be elements of $\mathcal{S}^\sharp$ that admit a common right-multiple, and let h be the right-lcm of s and t. By

assumption, we have $h = s(s\backslash t) = t(t\backslash s)$. As $\mathcal{S}^\sharp$ is closed under right-complement, there exist s', t' in $\mathcal{S}^\sharp$ and h' satisfying $st' = ts'$, $t\backslash s = s'h'$, and $s\backslash t = t'h'$, whence $h = (st')h'$. By definition of the right-lcm, h' must be an identity-element and, therefore, $s\backslash t$ and $t\backslash s$ belong to $\mathcal{S}^\sharp$. So $\mathcal{S}^\sharp$ is closed under the right-complement operation, and (i) implies (ii).

Conversely, assume that $\mathcal{S}^\sharp$ is closed under the right-complement operation. Assume that s, t belong to $\mathcal{S}^\sharp$ and $sg = tf$ holds. Then sg is a common right-multiple of s and t, hence it is a right-multiple of their right-lcm, which is $s(s\backslash t)$ and $t(t\backslash s)$. So there exists h satisfying $sg = s(s\backslash t)h$. Left-cancelling s, we deduce $g = (s\backslash t)h$ and, symmetrically, $f = (t\backslash s)h$. Then $t\backslash s$, $s\backslash t$, and h witness that the expected instance of closure under right-complement is satisfied. So (ii) implies (i). □

Solution to selected exercises

Exercice 38 (**multiplication by invertible**).— *Assume that $\mathcal{C}$ is a cancellative category, and $\mathcal{S}$ is a subfamily of $\mathcal{C}$ that is closed under left-divisor and contains at least one element with source x for each object x. Prove $\mathcal{S}^\sharp = \mathcal{S}$.*

Solution. Assume $g \in \mathcal{S}$ and $\epsilon \in \mathcal{C}^\times$. Then we have $g = g\epsilon\epsilon^{-1}$, whence $g\epsilon \preccurlyeq g$, and $g\epsilon \in \mathcal{S}$. So $\mathcal{S}^\sharp$ is included in $\mathcal{S}$.

Exercice 40 (**head** *vs.* **lcm**).— *Assume that $\mathcal{C}$ is a left-cancellative category, $\mathcal{S}$ is included in $\mathcal{C}$, and g belongs to $\mathcal{C} \setminus \mathcal{C}^\times$. Show that s is an $\mathcal{S}$-head of g if and only if it is a right-lcm of* $\mathrm{Div}(g) \cap \mathcal{S}$.

Solution. If s is an $\mathcal{S}$-head of g, then there exists g' so that $s|g'$ is an $\mathcal{S}$-greedy decomposition of g. By Lemma IV.1.21, we deduce that, for t in $\mathcal{S}$, the relation $t \preccurlyeq g$ implies $t \preccurlyeq s$: this means that every element of $\mathrm{Div}(g) \cap \mathcal{S}$ divides s and, therefore, that s is a left-lcm of $\mathrm{Div}(g) \cap \mathcal{S}$.

Exercice 41 (**closed under right-comultiple**).— *Assume that $\mathcal{C}$ is a left-cancellative category, $\mathcal{S}$ is a subfamily of $\mathcal{C}$, and there exists $H : \mathcal{C} \setminus \mathcal{C}^\times \to \mathcal{S}$ satisfying (IV.1.46). Show that $\mathcal{S}$ is closed under right-comultiple.*

Solution. Assume $f, g \in \mathcal{S}$ and $f\hat{g} = g\hat{f}$. If $f\hat{g}$ is invertible, then everything is obvious. Otherwise, $H(f\hat{g})$ is defined, and, by (IV.1.46)(i), we have $H(f\hat{g}) \preccurlyeq f\hat{g}$. On the other hand, we have $f \in \mathcal{S}$ and $f \preccurlyeq f\hat{g}$, whence $f \preccurlyeq H(f\hat{g})$ by (IV.1.46)(iii). A symmetric argument gives $g \preccurlyeq H(f\hat{g})$ as we have $f\hat{g} = g\hat{f}$. So $H(f\hat{g})$, which belongs to $\mathcal{S}$ by construction, is a common right-multiple of f and g of the expected type.

Exercice 42 (**power**).— *Assume that $\mathcal{S}$ is a Garside family a left-cancellative category $\mathcal{C}$. Show that, if $g_1|\cdots|g_p$ is an $\mathcal{S}^m$-normal decomposition of g and, for every i, the path $s_{i,1}|\cdots|s_{i,m}$ is an $\mathcal{S}$-normal decomposition of g_i, then the path $s_{1,1}|\cdots|s_{1,m}|s_{2,1}|\cdots|s_{2,m}|\cdots|s_{p,1}|\cdots|s_{p,m}$ is an $\mathcal{S}$-normal decomposition of g.*

Solution. By assumption, every $s_{i,j}$ lies in $\mathcal{S}^\sharp$ and $s_{i,j}|s_{i,j+1}$ is $\mathcal{S}$-greedy for all i, j. So the point is to show that $s_{i,m}|s_{i+1,1}$ is $\mathcal{S}$-greedy. Now assume $t \preccurlyeq s_{i,m}s_{i+1,1}$ with t in $\mathcal{S}$. Then we deduce $s_{i,1}\cdots s_{i,m-1}t \preccurlyeq s_{i,1}\cdots s_{i,m-1}s_{i,m}s_{i+1,1}$, whence $s_{i,1}\cdots s_{i,m-1}t \preccurlyeq g_ig_{i+1}$. As $g_i|g_{i+1}$ is $\mathcal{S}^m$-greedy, we deduce $s_{i,1}\cdots s_{i,m-1}t \preccurlyeq g_i$, that is, $s_{i,1}\cdots s_{i,m-1}t \preccurlyeq s_{i,1}\cdots s_{i,m-1}s_{i,m}$, which implies $t \preccurlyeq s_{i,m}$ since $\mathcal{C}$ is left-cancellative. As $\mathcal{S}$ is a Garside family, Corollary IV.1.31 implies that this is enough to conclude that $s_{i,m}|s_{i+1,1}$ is $\mathcal{S}$-greedy.

Exercice 45 (**no conditional right-lcm**).— *Let M be the monoid generated by $\mathtt{a}, \mathtt{b}, \mathtt{a}', \mathtt{b}'$ subject to the relations $\mathtt{ab}{=}\mathtt{ba}, \mathtt{a}'\mathtt{b}'{=}\mathtt{b}'\mathtt{a}', \mathtt{aa}'{=}\mathtt{bb}', \mathtt{a}'\mathtt{a}{=}\mathtt{b}'\mathtt{b}$.* (i) *Show that the cube condition is satisfied on $\{\mathtt{a}, \mathtt{b}, \mathtt{a}', \mathtt{b}'\}$, and that right- and left-reversing are complete for the above presentation.* (ii) *Show that M is cancellative and admits right-mcms.* (iii) *Show that $\mathtt{a}$ and $\mathtt{b}$ admit two right-mcms in M, but they admit no right-lcm.* (iv) *Let $S = \{\mathtt{a}, \mathtt{b}, \mathtt{a}', \mathtt{b}', \mathtt{ab}, \mathtt{a}'\mathtt{b}', \mathtt{aa}', \mathtt{a}'\mathtt{a}\}$, see diagram on the side. Show that S is closed under right-mcm, and deduce that $\mathcal{S}$ is a Garside family in M.* (v) *Show that the (unique) strict S-normal decomposition of the element $\mathtt{a}^2\mathtt{b}'\mathtt{a}'^2$ is $\mathtt{ab}|\mathtt{a}'\mathtt{b}'|\mathtt{b}'$.*



Solution. (i) There are several relations of the form $\mathtt{a}\cdots = \mathtt{b}\cdots$, but, as there is no relation $\mathtt{a}\cdots = \mathtt{a}'\cdots$, there is no mixed cube to consider: the only triples to check are $(\mathtt{a}, \mathtt{a}, \mathtt{b})$ and the like, and this is easy.

(ii) The presentation is homogeneous, hence (strongly) Noetherian. By Lemma II.4.62, right-reversing is complete. Hence M is right-cancellative and admits right-mcms. By symmetry of the relations, left-reversing is complete as well, and M is right-cancellative.

(iii) By assumption, $\mathtt{ab}$ and $\mathtt{aa}'$ are common right-multiples of $\mathtt{a}$ and $\mathtt{b}$. Owing to their length, they must be minimal. As right-reversing is complete, every common right-multiple of $\mathtt{a}$ and $\mathtt{b}$ is a right-multiple or $\mathtt{ab}$ and $\mathtt{aa}'$. Hence the latter are the only mcms of $\mathtt{a}$ and $\mathtt{b}$. Finally neither is a right-multiple of the other, so $\mathtt{a}$ and $\mathtt{b}$ admit no right-lcm.

(iv) The set S generates M, and it is closed under right-divisor and right-mcm: $\mathtt{ab}$ and $\mathtt{aa}'$ are common right-multiples of $\mathtt{a}$ and $\mathtt{b}$, $\mathtt{a}'\mathtt{b}'$ and $\mathtt{a}'\mathtt{a}$ are the two right-mcms of $\mathtt{a}'$ and $\mathtt{b}'$, whereas none of the pairs $\{\mathtt{a}, \mathtt{a}'\}$, $\{\mathtt{a}, \mathtt{b}'\}$, $\{\mathtt{b}, \mathtt{b}'\}$, and $\{\mathtt{b}, \mathtt{a}'\}$ admits a common right-multiple. Hence, by Corollary IV.2.26, S is a Garside family in M.

(v) Push the letters to the left as much as possible.

Exercice 47 (**solid**).— *Assume that $\mathcal{C}$ is a left-cancellative category and $\mathcal{S}$ is a generating subfamily of $\mathcal{C}$.* (i) *Show that $\mathcal{S}$ is solid in $\mathcal{C}$ if and only if $\mathcal{S}$ includes $\mathbf{1}_{\mathcal{C}}$ and it is closed under right-quotient.* (ii) *Assume moreover that $\mathcal{S}$ is closed under right-divisor. Show that $\mathcal{S}$ includes $\mathcal{C}^\times \setminus \mathbf{1}_{\mathcal{C}}$, that $\epsilon \in \mathcal{S} \cap \mathcal{C}^\times$ implies $\epsilon^{-1} \in \mathcal{S}$, and that and $\mathcal{C}^\times\mathcal{S} = \mathcal{S}$ holds, but that that $\mathcal{S}$ need not be solid.*

Solution. (ii) Assume that ϵ is a nontrivial invertible element, with target y. As $\mathcal{S}$ generates $\mathcal{C}$, there exists at least one element ϵ' of $\mathcal{S}$ that right-divides ϵ. Then 1_y right-divides ϵ', which lies in $\mathcal{S}$, hence 1_y must lie in $\mathcal{S}$. Next, ϵ right-divides 1_y, which lies in $\mathcal{S}$, hence ϵ must lie in $\mathcal{S}$. Now assume $\epsilon \in \mathcal{S} \cap \mathcal{C}^\times$. Then either ϵ is an identity-element, in which case it coincides with ϵ^{-1} and the latter lies in $\mathcal{S}$, or ϵ^{-1} is invertible and is not an identity-element, in which case it belongs to $\mathcal{S}$ by the above argument. For $\mathcal{C}^\times\mathcal{S} = \mathcal{S}$, the proof is as for Lemma IV.2.2. If y is an object

that is the target of no non-identity-element, then 1_y: need not belong to $\mathcal{S}$. For instance, in the monoid $\{1\}$, the empty set satisfies the condition.

Exercice 48 (**solid**).— *Let M be the monoid $\langle \mathtt{a}, \mathtt{e} \,|\, \mathtt{ea} = \mathtt{a}, \mathtt{e}^2 = 1\rangle^+$.* (i) *Show that every element of M has a unique expression of the form $\mathtt{a}^p\mathtt{e}^q$ with $p \geqslant 0$ and $q \in \{0, 1\}$, and that M is left-cancellative.* (ii) *Let $S = \{1, \mathtt{a}, \mathtt{e}\}$. Show that S is a solid Garside family in M, but that $S = S^\sharp$ does not hold.*

Solution. (i) Every word in $\{\mathtt{a}, \mathtt{e}\}^*$ is equivalent to a word of the form $\mathtt{a}^p$ or $\mathtt{a}^p\mathtt{e}$. Conversely, it is impossible that two distinct words of this form are equivalent, as the number of $\mathtt{a}$ is an invariant, and so is the parity of the number of $\mathtt{e}$ that follows the last $\mathtt{a}$. The formulas $\mathtt{a} \cdot \mathtt{a}^p\mathtt{e}^q = \mathtt{a}^{p+1}\mathtt{e}^q$ and $\mathtt{e} \cdot \mathtt{a}^p\mathtt{e}^q = \mathtt{a}^p\mathtt{e}^q$ for $p \geqslant 1$, plus $\mathtt{e} \cdot \mathtt{e}^q = \mathtt{e}^{q+1 \mod 2}$ show that, for every s in $\{\mathtt{a}, \mathtt{e}\}$, the value of g can be recovered from s and sg. (ii) S generates M. The explicit formula for the multiplication shows that $\mathtt{a}$ is right-divisible only by 1 and itself, and so does $\mathtt{e}$. Hence S is closed under right-divisor, and it is solid. For $p \geqslant 1$, define $H(\mathtt{a}^p\mathtt{e}^q) = a$. Then H is a S-head function. Hence, by Proposition IV.2.7(i), S is a Garside family in M. On the other hand, $\mathtt{ae}$ is an element of $S^\sharp \setminus S$.

Exercice 49 (**not solid**).— *Let $M = \langle \mathtt{a}, \mathtt{e} \,|\, \mathtt{ea} = \mathtt{ae}, \mathtt{e}^2 = 1\rangle^+$, and $S = \{\mathtt{a}, \mathtt{e}\}$.* (i) *Show that M is left-cancellative. [Hint: M is $\mathbb{N} \times \mathbb{Z}/2\mathbb{Z}$]* (ii) *Show that S is a Garside family of M, but S is not solid in M. [Hint: $\mathtt{ea}$ right-divides $\mathtt{a}$, but does not belong to S.]*

Solution. (i) Every word of $\{\mathtt{a}, \mathtt{e}\}^*$ is equivalent modulo the relations to a word of the form $\mathtt{a}^p\mathtt{e}^q$ with $q \leqslant 1$. As $(1, \dot{0})$ and $(0, \dot{1})$ satisfy in $\mathbb{N}\times\mathbb{Z}/2\mathbb{Z}$ the defining relations of M, there exists a well defined homomorphism F of M to $\mathbb{N}\times\mathbb{Z}/2\mathbb{Z}$ that satisfies $F(\mathtt{a}) = (1, \dot{0})$ and $F(\mathtt{e}) = (0, \dot{1})$. As $(1, \dot{0})$ and $(0, \dot{1})$ generate $\mathbb{N}\times\mathbb{Z}/2\mathbb{Z}$, the homomorphism F is surjective. As the images under F of pairwise distinct words $\mathtt{a}^p\mathtt{e}^q$ are distinct, F is injective, hence it is a isomorphism. Hence $\mathbb{N}\times\mathbb{Z}/2\mathbb{Z}$ is left-cancellative, so is M. (ii) By definition, S generates M. Next, the invertible elements of M are 1 and $\mathtt{e}$. The equality $\mathtt{ea} = \mathtt{ae}$ implies $M^\times S \subseteq S^\sharp$. Finally, the three elements of S^2, namely $\mathtt{a}^2, \mathtt{ae}$, and 1, admit S-normal decompositions, namely for instance $\mathtt{a}|\mathtt{a}$, $\mathtt{a}|\mathtt{e}$, and ε. So S is a Garside family in M. Now $\mathtt{ae}$ does not lie in S, but we have $\mathtt{a} = \mathtt{e}(\mathtt{ae})$, so S is not closed under right-divisor, hence is not solid.

Exercice 50 (**recognizing Garside, right-lcm solid case**).— *Assume that $\mathcal{S}$ is a solid subfamily in a left-cancellative category $\mathcal{C}$ that is right-Noetherian and admits conditional right-lcms. Show that $\mathcal{S}$ is a Garside family in $\mathcal{C}$ if and only if $\mathcal{S}$ generates $\mathcal{C}$ and it is weakly closed under right-lcm.*

Solution. If $\mathcal{S}$ is a (solid or not solid) Garside family in $\mathcal{C}$, then, by Corollary IV.2.29, $\mathcal{S}$ is weakly closed under right-lcm. Conversely, assume that $\mathcal{S}$ is solid, generates $\mathcal{C}$, and is weakly closed under right-lcm. By definition, $\mathcal{S}$ is closed under right-divisor, hence, by Lemma IV.1.13(i), so is $\mathcal{S}^\sharp$. Then Corollary IV.2.29(iii) implies that $\mathcal{S}$ is a Garside family in $\mathcal{C}$.

Exercice 52 (**local right-divisibility**).— *Assume that $\mathcal{C}$ is a left-cancellative category and $\mathcal{S}$ is a generating subfamily of $\mathcal{C}$ that is closed under right-divisor.* (i) *Show that the transitive closure of $\widetilde{\preccurlyeq}_{\mathcal{S}}$ is the restriction of $\widetilde{\preccurlyeq}$ to $\mathcal{S}$.* (ii) *Show that*

the transitive closure of $\widetilde{\preccurlyeq}_{\mathcal{S}}$ is almost the restriction of $\widetilde{\preccurlyeq}$ to $\mathcal{S}$, in the following sense: if $s \widetilde{\preccurlyeq} t$ holds, there exists $s' \,{}^{\times}\!= s$ satisfying $s' \widetilde{\preccurlyeq}_{\mathcal{S}}^{*} t$.

Solution. Let $\widetilde{\preccurlyeq}_{\mathcal{S}}^{*}$ be the transitive closure of $\widetilde{\preccurlyeq}_{\mathcal{S}}$. As $s \widetilde{\preccurlyeq}_{\mathcal{S}} t$ implies $s \widetilde{\preccurlyeq} t$ trivially and $\widetilde{\preccurlyeq}$ is transitive, $s \widetilde{\preccurlyeq}_{\mathcal{S}}^{*} t$ implies $s \widetilde{\preccurlyeq} t$. Conversely, assume $s \widetilde{\preccurlyeq} t$, say $t = gs$. As $\mathcal{S}$ generates $\mathcal{C}$, there exist $s_1, \dots, s_q$ in $\mathcal{S}$ satisfying $g = s_1 \cdots s_p$. So we have $t = s_1 \cdots s_p s$. Put $t_i = s_i \cdots s_p s$ for $1 \leqslant i \leqslant p$. Each element t_i right-divides t, an element of $\mathcal{S}$, so it belongs to $\mathcal{S}$. Now, by construction, we have $s = \widetilde{\preccurlyeq}_{\mathcal{S}} t_1 \widetilde{\preccurlyeq}_{\mathcal{S}} \cdots \widetilde{\preccurlyeq}_{\mathcal{S}} t_p = t$, whence $s \widetilde{\preccurlyeq}_{\mathcal{S}}^{*} t$.

(ii) Let $\widetilde{\preccurlyeq}_{\mathcal{S}}^{*}$ be the transitive closure of $\widetilde{\preccurlyeq}_{\mathcal{S}}$. By Lemma IV.2.15, $s \widetilde{\preccurlyeq}_{\mathcal{S}} t$ implies $s \widetilde{\preccurlyeq} t$, hence $s \widetilde{\preccurlyeq}_{\mathcal{S}}^{*} t$ implies $s \widetilde{\preccurlyeq} t$ since $\widetilde{\preccurlyeq}$ is transitive. Conversely, assume $s \widetilde{\preccurlyeq} t$, say $t = gs$ with $g \notin \mathcal{C}^{\times}$. As $\mathcal{S}$ generates $\mathcal{C}$, there exist $s_1, \dots, s_q$ in $\mathcal{S}$ satisfying $g = s_1 \cdots s_p$. Assume that p has been chosen minimal. Then $s_1, \dots, s_{p-1}$ are not invertible: if s_i is invertible, then $s_i s_{i+1}$ right-divides s_{i+1}, hence belongs to $\mathcal{S}$, and therefore grouping s_i and s_{i+1} would provide a shorter decomposition. We have $t = s_1 \cdots s_p s$. Put $t_i = s_i \cdots s_p s$ for $1 \leqslant i \leqslant p$. Each element t_i right-divides t, an element of $\mathcal{S}$, so it belongs to $\mathcal{S}$. As $\mathcal{S}$ is closed under right-divisor, we have $\mathcal{C}^{\times}\mathcal{S} \subseteq \mathcal{S}$. So we can assume that $s_1, \dots, s_{p-1}$ are non-invertible. Hence we have $s_1 s \widetilde{\preccurlyeq}_{\mathcal{S}}^{*} t$.

Exercice 53 (**local left-divisibility**).— *Assume that $\mathcal{S}$ is a subfamily of a left-cancellative category $\mathcal{C}$.* (i) *Show that $s \preccurlyeq_{\mathcal{S}} t$ implies $s \preccurlyeq t$, and that $s \preccurlyeq_{\mathcal{S}} t$ is equivalent to $s \preccurlyeq t$ whenever $\mathcal{S}$ is closed under right-quotient in $\mathcal{C}$.* (ii) *Show that, if $\mathcal{S}^{\times} = \mathcal{C}^{\times} \cap \mathcal{S}$ holds, then $s \prec_{\mathcal{S}} t$ implies $s \prec t$.* (iii) *Show that, if $\mathcal{S}$ is closed under right-divisor, then $\preccurlyeq_{\mathcal{S}}$ is the restriction of $\preccurlyeq$ to $\mathcal{S}$ and, if $\mathcal{S}^{\times} = \mathcal{C}^{\times} \cap \mathcal{S}$ holds, $\prec_{\mathcal{S}}$ is the restriction of $\prec$ to $\mathcal{S}$.*

Solution. (iii) First $s \preccurlyeq_{\mathcal{S}} t$ implies $s \preccurlyeq t$. Conversely, assume $s, t \in \mathcal{S}$ and $sg' = t$. As t belongs to $\mathcal{S}$, the assumption that $\mathcal{S}$ is closed under right-divisor implies that g' belongs to $\mathcal{S}$, hence witnesses for $s \preccurlyeq_{\mathcal{S}} t$. Next, by Lemma IV.2.15, $s \prec_{\mathcal{S}} t$ implies $s \prec t$. Conversely, assume $st' = t$ with $t' \notin \mathcal{C}^{\times}$. As above, t' must belong to $\mathcal{S}$, and it cannot belong to $\mathcal{S}^{\times}$. So $s \prec_{\mathcal{S}} t$ holds.

Exercice 55 (**locally right-Noetherian**).— *Assume that $\mathcal{C}$ is a left-cancellative category and $\mathcal{S}$ is a subfamily of $\mathcal{C}$.* (i) *Prove that $\mathcal{S}$ is locally right-Noetherian if and only if, for every s in $\mathcal{S}^{\sharp}$, every $\prec_{\mathcal{S}}$-increasing sequence in $\mathrm{Div}_{\mathcal{S}}(s)$ is finite.* (ii) *Assume that $\mathcal{S}$ is locally right-Noetherian and closed under right-divisor. Show that $\mathcal{S}^{\sharp}$ is locally right-Noetherian. [Hint: For $\mathcal{X} \subseteq \mathcal{S}^{\sharp}$ introduce $\mathcal{X}' = \{s \in \mathcal{S} \mid \exists \epsilon, \epsilon' \in \mathcal{C}^{\times}\, (\epsilon s \epsilon' \in \mathcal{X})\}$, and construct a $\widetilde{\preccurlyeq}$-minimal element in $\mathcal{X}$ from a $\widetilde{\preccurlyeq}$-minimal element in $\mathcal{X}'$.]*

Solution. (ii) Let $\mathcal{X}$ be a nonempty subfamily of $\mathcal{S}^{\sharp}$. Put

$$\mathcal{X}' = \{s \in \mathcal{S} \mid \exists \epsilon, \epsilon' \in \mathcal{C}^{\times}\, (\epsilon s \epsilon' \in \mathcal{X})\}.$$

We have $\mathcal{X} \subseteq \mathcal{X}'$, whence $\mathcal{X}' \neq \emptyset$. As $\mathcal{X}'$ is included in $\mathcal{S}$, it contains a $\widetilde{\preccurlyeq}_{\mathcal{S}}$-minimal element, say s_0. By definition, there exists ϵ_0, ϵ_0' invertible such that $\epsilon_0 s_0 \epsilon_0'$ lies in $\mathcal{X}$. Put $t_0 = \epsilon_0 s_0 \epsilon_0'$. We claim that t_0 is $\widetilde{\preccurlyeq}_{\mathcal{S}^{\sharp}}$-minimal in $\mathcal{X}$. Indeed, assume $t_0 = rt$ with r in $\mathcal{S}^{\sharp}$ and t in $\mathcal{X}$. We have to prove that r lies in $(\mathcal{S}^{\sharp})^{\times}$. As $\mathcal{S}^{\sharp}$ is solid, it suffices to show that r lies in $\mathcal{C}^{\times}$. By definition, r belongs to $\mathcal{S}\mathcal{C}^{\times} \cup \mathcal{C}^{\times}$. If r belongs to $\mathcal{C}^{\times}$, we are done. Otherwise, write $r = r'\epsilon$ with r' in $\mathcal{S}$ and ϵ in $\mathcal{C}^{\times}$. Then we have $s_0 = (\epsilon_0^{-1} r')(\epsilon t \epsilon_0'^{-1})$. As s_0 lies in $\mathcal{S}$ and $\mathcal{S}$ is closed under right-divisor,

$\epsilon t {\epsilon'_0}^{-1}$ lies in $\mathcal{S}$. As t belongs to $\mathcal{X}$, we deduce that $\epsilon t {\epsilon'_0}^{-1}$ lies in $\mathcal{X}'$. Then, by the choice of s_0, the elements $\epsilon_0^{-1} r'$, and therefore r' and $r'\epsilon$, that is, r, must be invertible in $\mathcal{C}$. Hence t_0 is $\widetilde{\preccurlyeq}_{\mathcal{S}^\sharp}$-minimal in $\mathcal{X}$, and $\widetilde{\preccurlyeq}_{\mathcal{S}^\sharp}$ is a well-founded relation, that is, $\mathcal{S}^\sharp$ is locally right-Noetherian.

Chapter V: Bounded Garside families

Skipped proofs

Proposition V.1.59 (**right-cancellative II**).— *If $\mathcal{S}$ is a Garside family in a left-cancellative category $\mathcal{C}$ that is right-bounded by a target-injective map Δ, and ϕ_Δ preserves $\mathcal{S}$-normality and is surjective on $\mathcal{C}^\times$, then the following conditions are equivalent:*

(i) *The category $\mathcal{C}$ is right-cancellative;*
(ii) *The functor ϕ_Δ is injective on $\mathcal{C}$;*
(iii) *The functor ϕ_Δ is injective on $\mathcal{S}^\sharp$.*

Proof. By Proposition V.1.36, (i) and (ii) are equivalent, and (ii) obviously implies (iii).

Now assume that ϕ_Δ is injective on $\mathcal{S}^\sharp$. We will prove that it is injective on $\mathcal{C}$. First, we claim that $\phi_\Delta(s') =^\times \phi_\Delta(s)$ implies $s =^\times s'$ for s, s' in $\mathcal{S}^\sharp$. Indeed, assume $\phi_\Delta(s') = \phi_\Delta(s)\,\epsilon$ with $\epsilon \in \mathcal{C}^\times$. If ϵ is trivial, the injectivity of ϕ_Δ on $\mathcal{S}^\sharp$ implies $s' = s$. Otherwise, as ϕ_Δ is surjective on $\mathcal{C}^\times$, there exists ϵ' in $\mathcal{C}^\times$ satisfying $\phi_\Delta(\epsilon') = \epsilon$. As Δ is target-injective, the assumption that $\phi_\Delta(s)\phi_\Delta(\epsilon')$ is defined implies that $s\epsilon'$ is defined too: if y is the target of s and x' is the source of ϵ', we obtain $\phi_\Delta(y) = \phi_\Delta(x')$, whence $y = x'$ as ϕ_Δ is injective on objects. We deduce $\phi_\Delta(s') = \phi_\Delta(s\epsilon')$. As s belongs to $\mathcal{S}^\sharp$, so does $s\epsilon'$, and the assumption that ϕ_Δ is injective on $\mathcal{S}^\sharp$ then implies $s' = s\epsilon'$, hence $s' =^\times s$.

Now we prove using induction on $p \geqslant 1$ the statement: $\phi_\Delta(g) = \phi_\Delta(g')$ implies $g = g'$ for all g, g' satisfying $\max(\|g\|_\mathcal{S}, \|g'\|_\mathcal{S}) \leqslant p$. For $p = 1$, as $\|g\|_\mathcal{S} \leqslant 1$ implies $g \in \mathcal{S}^\sharp$, the result follows from the assumption that ϕ_Δ is injective on $\mathcal{S}^\sharp$. Assume $p \geqslant 2$, and $\phi_\Delta(g) = \phi_\Delta(g')$ with $\max(\|g\|_\mathcal{S}, \|g'\|_\mathcal{S}) \leqslant p$. Let $s_1|\cdots|s_p$ and $s'_1|\cdots|s'_p$ be $\mathcal{S}$-normal decompositions of g and g'. As ϕ_Δ is a functor and it preserves normality, $\phi_\Delta(s_1)|\cdots|\phi_\Delta(s_p)$ and $\phi_\Delta(s'_1)|\cdots|\phi_\Delta(s'_p)$ are $\mathcal{S}$-normal decompositions of $\phi_\Delta(g)$ and $\phi_\Delta(g')$. By Proposition III.1.25 (normal unique), $\phi_\Delta(g) = \phi_\Delta(g')$ implies $\phi_\Delta(s_1) =^\times \phi_\Delta(s'_1)$, whence, by the claim above, $s'_1 =^\times s_1$. Hence s_1 is an $\mathcal{S}$-head for g and g', and we can write $g = s_1 g_1$, $g' = s_1 g'_1$, with $\max(\|g_1\|_\mathcal{S}, \|g'_1\|_\mathcal{S}) \leqslant p-1$. Then $\phi_\Delta(g') = \phi_\Delta(g)$ implies $\phi_\Delta(s_1)\phi_\Delta(g_1) = \phi_\Delta(s_1)\phi_\Delta(g'_1)$, whence $\phi_\Delta(g_1) = \phi_\Delta(g'_1)$. The induction hypothesis implies $g_1 = g'_1$, whence $g' = g$. So ϕ_Δ is injective on $\mathcal{C}$, and (iii) implies (ii). □

Proposition V.2.34 (**right-cancellativity III**).— *If $\mathcal{C}$ is a left-cancellative category and Δ is a Garside map of $\mathcal{C}$ that preserves Δ-normality, then $\mathcal{C}$ is right-cancellative if and only if ϕ_Δ is injective on $(\mathcal{D}iv(\Delta))^2$.*

Proof. If $\mathcal{C}$ is right-cancellative, Proposition V.1.36 implies that ϕ_Δ is injective on $\mathcal{C}$, hence a fortiori on $(\mathcal{D}iv(\Delta))^2$, so the condition is necessary.

Conversely, assume that ϕ_Δ is injective on $(\mathcal{D}iv(\Delta))^2$. First, ϕ_Δ must be injective on $\mathbf{1}_\mathcal{C}$, which is included in $(\mathcal{D}iv(\Delta))^2$, and, therefore, on $\mathcal{O}bj(\mathcal{C})$, that is, Δ must be target-injective. Next, as $\mathbf{1}_\mathcal{C}$ is included in $\mathcal{D}iv(\Delta)$, the assumption implies that ϕ_Δ is injective on $\mathcal{D}iv(\Delta)$.

We claim that ϕ_Δ induces a permutation of $\mathcal{C}^\times$. Indeed, as ϕ_Δ is a functor, it maps $\mathcal{C}^\times$ into itself. Now, the assumption that Δ is a Garside map implies that $\mathcal{D}iv(\Delta)$ is bounded by Δ, hence, by Lemma V.2.7, ϕ_Δ induces a surjective map of $\mathcal{D}iv(\Delta)$ into itself, hence a permutation of $\mathcal{D}iv(\Delta)$ as it is injective on $\mathcal{D}iv(\Delta)$. Assume $\epsilon \in \mathcal{C}^\times$. Then ϵ and ϵ^{-1} belong to $\mathcal{D}iv(\Delta)$, so there exist s, t in $\mathcal{D}iv(\Delta)$ satisfying $\phi_\Delta(s) = \epsilon$ and $\phi_\Delta(t) = \epsilon^{-1}$. As Δ is target-injective, st is defined. Indeed, let y be the target of s and x' be the source of t. As ϕ_Δ is a functor, $\phi_\Delta(y)$ is the target of $\phi_\Delta(s)$, that is, of ϵ, whereas $\phi_\Delta(x')$ is the source of $\phi_\Delta(t)$, that is, of ϵ^{-1}. Hence we have $\phi_\Delta(y) = \phi_\Delta(x')$ and, therefore, $y = x'$, that is, st is defined. The argument for ts is symmetric. So st and ts belong to $(\mathcal{D}iv(\Delta))^2$. The assumption that ϕ_Δ is injective on $(\mathcal{D}iv(\Delta))^2$ implies $st = 1_x$ and $ts = 1_y$, where x (*resp.* y) is the source (*resp.* target) of s. So s belongs to $\mathcal{C}^\times$, and ϕ_Δ induces a permutation of $\mathcal{C}^\times$. Then, Proposition V.1.59 implies that $\mathcal{C}$ is right-cancellative.

Note that the assumptions of Proposition V.2.34 can be slightly weakened: the only assumptions used in the proof is that ϕ_Δ is injective on $\mathcal{D}iv(\Delta)$ and that, for g in $(\mathcal{D}iv(\Delta))^2$, the relation $\phi_\Delta(g) \in \mathbf{1}_\mathcal{C}$ implies $g \in \mathbf{1}_\mathcal{C}$. We do not know whether the latter condition can be skipped. □

Lemma V.2.38.— (i) *A left-cancellative category that is left-Noetherian and admits left-gcds admits conditional right-lcms.*

(ii) *In a cancellative category that admits conditional right-lcms, any two elements of $\mathcal{C}$ that admit a common left-multiple admit a right-gcd.*

Proof. (i) We first show that every nonempty family of elements of $\mathcal{C}$ sharing the same source admits a left-gcd. Let $\mathcal{S}$ be a nonempty family of elements of $\mathcal{C}$ that share the same source. An obvious induction shows that, in $\mathcal{C}$, every finite nonempty family of elements of $\mathcal{C}$ sharing the same source has a left-gcd. For Y a finite nonempty subset of $\mathcal{S}$, choose a left-gcd g_Y for Y, and let $\mathcal{S}'$ be the family of all elements g_Y. As $\mathcal{C}$ is left-Noetherian, there exists an element g_X of $\mathcal{S}'$ that is $\prec$-minimal in $\mathcal{S}'$. We claim that g_X is a left-gcd for $\mathcal{S}$. Indeed, let g be an arbitrary element of $\mathcal{S}$. The point is to prove $g_X \preccurlyeq g$. Now, by construction, we have $g_{X\cup\{g\}} \preccurlyeq g$ and $g_{X\cup\{g\}} \preccurlyeq g_X$. As g_X is $\prec$-minimal in $\mathcal{S}'$, we must have $g_{X\cup\{g\}} =^\times g_X$, whence $g_X \preccurlyeq g_{X\cup\{g\}} \preccurlyeq g$, as expected. So $\mathcal{S}$ has a left-gcd. Then Lemma II.2.21 guarantees that $\mathcal{C}$ admits conditional right-lcms.

(ii) (See Figure 8.) Let f, g be two elements of $\mathcal{C}$ that admit a common left-multiple, say $f'g = g'f$, hence share the same target. The elements f' and g' admit a common right-multiple, namely $f'g$, hence they admit a right-lcm, say $f'g'' = g'f''$. By definition of a right-lcm, there exists h satisfying $f = f''h$ and $g = g''h$. By construction, h is a common right-divisor of f and g.

Let $\hat{h}$ be a common right-divisor of f and g. So there exist $\hat{f}, \hat{g}$ satisfying $f = \hat{f}\hat{h}$ and $g = \hat{g}\hat{h}$. Then we have $f'\hat{g}\hat{h} = f'g = g'f = g'\hat{f}\hat{h}$, whence $f'\hat{g} = g'\hat{f}$ by right-cancelling $\hat{h}$. So $f'\hat{g}$ is a common right-multiple of f' and g', hence it is a

right-multiple of their right-lcm, that is, there exists h' satisfying $f'\hat{g} = (f'g'')h'$, whence $\hat{g} = g''h'$ by left-cancelling f'. We deduce $g''h = g = \hat{g}\hat{h} = g''h'\hat{h}$, whence $h = h'\hat{h}$ by right-cancelling $\hat{h}$. So $\hat{h}$ right-divides h, which shows that h is a right-gcd of f and g. □



FIGURE 8. Construction of a right-gcd using a right-lcm.

Lemma V.3.40.— *If Δ is a Garside map in a cancellative category $\mathcal{C}$, then, for all s, t in $\mathcal{D}iv(\Delta)$, we have $s \preccurlyeq t \Leftrightarrow \partial s \succcurlyeq \partial t$ and $s \succcurlyeq t \Leftrightarrow \widetilde{\partial} s \preccurlyeq \widetilde{\partial} t$.*

Proof. As ∂ and $\widetilde{\partial}$ are bijective, everything is clear. For instance, assume $s \preccurlyeq t$. Then we have $st' = t$ for some t', which necessarily belongs to $\mathcal{D}iv(\Delta)$. Calling x the common source of s and t, we deduce

$$\Delta(x) = s\,\partial s = t\,\partial t = s\,t'\,\partial t,$$

whence $\partial s = t'\,\partial t$, and $\partial s \succcurlyeq \partial t$. The other implications are similar. □

Lemma V.3.46.— *If Δ is a Garside map in a cancellative category $\mathcal{C}$ and f, g are elements of $\mathcal{C}(x, \text{-})$, then, for every m satisfying $m \geqslant \max(\sup_\Delta(f), \sup_\Delta(g))$ and for every h in $\mathrm{Div}(\Delta^{[m]}(x))$, the following conditions are equivalent:*

(i) *The element h is a right-lcm of f and g;*

(ii) *The element $\partial^{[m]}h$ is a right-gcd of $\partial^{[m]}f$ and $\partial^{[m]}g$.*

Proof. Assume first that $m = 1$. As $\mathcal{D}iv(\Delta)$ is closed under right-comultiple, h is a right-lcm of f and g if and only if the three relations

$$f \preccurlyeq h,\ g \preccurlyeq h,\ \text{and } \forall h' \in \mathcal{D}iv(\Delta)\ ((f \preccurlyeq h' \text{ and } g \preccurlyeq h') \text{ implies } h \preccurlyeq h')$$

are satisfied. By the duality formulas of Lemma V.3.40, this is equivalent to

$$\partial f \succcurlyeq \partial h,\ \partial g \succcurlyeq \partial h,\ \text{and } \forall h' \in \widetilde{\mathcal{D}iv}(\Delta)\ ((\partial f \succcurlyeq h' \text{ and } \partial g \succcurlyeq h') \text{ implies } \partial h \succcurlyeq h'),$$

hence, as $\widetilde{\mathcal{D}iv}(\Delta)$ coincides with $\mathcal{D}iv(\Delta)$, to ∂h being a right-gcd of ∂f and ∂g.

Assume now that m is arbitrary. By Proposition V.2.32(iv), $\Delta^{[m]}$ is a Garside map in $\mathcal{C}$, and $\partial^{[m]}$ is the associated duality map. Applying the above result to $\Delta^{[m]}$ and $\partial^{[m]}$ gives the expected equivalence.

(Observe that, in the situation of Lemma V.3.46, $\Delta^{[m]}(x)$ is a common right-multiple of f and g, so every possible right-lcm of f and g must lie in $\mathrm{Div}(\Delta^{[m]}(x))$.) □

Solution to selected exercises

Exercice 59 (**preserving** $\mathcal{D}iv(\Delta)$).— *Assume that $\mathcal{C}$ is a category, Δ is a map from $\mathcal{O}bj(\mathcal{C})$ to $\mathcal{C}$ and ϕ is a functor from $\mathcal{C}$ into itself that commutes with Δ. Show that ϕ maps $\mathcal{D}iv(\Delta)$ and $\widetilde{\mathcal{D}iv}(\Delta)$ to themselves.*

Solution. Assume $s \in \mathcal{D}iv(\Delta)$, say $s \preccurlyeq \Delta(x)$. By (the easy direction of) Lemma II.2.8, this implies $\phi(s) \preccurlyeq \phi(\Delta(x))$. By assumption, the latter is $\Delta(\phi(x))$, so $\phi(s)$ belongs to $\mathcal{D}iv(\Delta)$. The argument is similar for $\widetilde{\mathcal{D}iv}(\Delta)$.

Exercice 60 (**preserving normality I**).— *Assume that $\mathcal{C}$ is a cancellative category, $\mathcal{S}$ is a Garside family of $\mathcal{C}$, and ϕ is a functor from $\mathcal{C}$ to itself.* (i) *Show that, if ϕ induces a permutation of $\mathcal{S}^\sharp$, then ϕ preserves $\mathcal{S}$-normality.* (ii) *Show that ϕ preserves non-invertibility, that is, $\phi(g)$ is invertible if and only if g is.*

Solution. (i) Assume that $s_1|s_2$ is $\mathcal{S}$-normal. First, by assumption, ϕ maps $\mathcal{S}^\sharp$ to itself, hence $\phi(s_1)$ and $\phi(s_2)$ lie in $\mathcal{S}^\sharp$. Assume that s is an element of $\mathcal{S}$ that satisfies $s \preccurlyeq \phi(s_1)\,\phi(s_2)$. By Lemma II.2.8, we deduce $\phi^{-1}(s) \preccurlyeq s_1 s_2$, whence $\phi^{-1}(s) \preccurlyeq s_1$ as ϕ^{-1} maps $\mathcal{S}^\sharp$ into itself and $s_1|s_2$, which is $\mathcal{S}$-normal by assumption, is also $\mathcal{S}^\sharp$-normal by Lemma III.1.10. Reapplying ϕ, we deduce $s \preccurlyeq \phi(s_1)$, and we conclude that $\phi(s_1)|\phi(s_2)$ is $\mathcal{S}$-normal.

(ii) First, $\phi(g)$ is invertible whenever g is invertible since ϕ is a functor. Conversely, assume that $\phi(g)$ is invertible. Let $g_1|\cdots|g_p$ be an $\mathcal{S}^\sharp$-normal decomposition of g. As ϕ is a functor and preserves normality, $\phi(g_1)|\cdots|\phi(g_p)$ is an $\mathcal{S}^\sharp$-normal decomposition of $\phi(g)$. The assumption that $\phi(g)$ is invertible implies that each of $\phi(g_1), \dots, \phi(g_p)$ is invertible, and then the assumption that ϕ is injective on $\mathcal{S}^\sharp$ implies that $g_1, \cdots, g_p$ are invertible. Hence g is invertible.

Exercice 61 (**preserving normality II**).— *Assume that $\mathcal{C}$ is a left-cancellative category and $\mathcal{S}$ is a Garside family of $\mathcal{C}$ that is right-bounded by a map Δ.* (i) *Show that ϕ_Δ preserves normality if and only if there exists an $\mathcal{S}$-head map H satisfying $H(\phi_\Delta(g)) =^\times \phi_\Delta(H(g))$ for every g in $(\mathcal{S}^\sharp)^2$, if and only if, for each $\mathcal{S}$-head map H, the above relation is satisfied.* (ii) *Show that a sufficient condition for ϕ_Δ to preserve normality is that ϕ_Δ preserves left-gcds on $\mathcal{S}^\sharp$, that is, if r, s, t belong to $\mathcal{S}^\sharp$ and r is a left-gcd of s and t, then $\phi_\Delta(r)$ is a left-gcd of $\phi_\Delta(s)$ and $\phi_\Delta(t)$.*

Solution. (i) Assume that ϕ_Δ preserves normality, H is a $\mathcal{S}$-head map, and $g_1|g_2$ lies in $(\mathcal{S}^\sharp)^{[2]}$. Put $g_1' = H(g_1 g_2)$, and let g_2' satisfy $g_1' g_2' = g_1 g_2$. Then $g_1'|g_2'$ is $\mathcal{S}$-normal, hence, as ϕ_Δ preserves normality, so is $\phi_\Delta(g_1')|\phi_\Delta(g_2')$. Moreover, we have $\phi_\Delta(g_1')\phi_\Delta(g_2') = \phi_\Delta(g_1 g_2)$, hence $\phi_\Delta(g_1')|\phi_\Delta(g_2')$ is an $\mathcal{S}$-normal decomposition of $\phi_\Delta(g_1 g_2)$. By uniqueness of the head, we must have $H(\phi_\Delta(g_1 g_2)) =^\times \phi_\Delta(g_1')$. Conversely, assume that H is a $\mathcal{S}^\sharp$-head map and $H(\phi_\Delta(g)) =^\times \phi_\Delta(H(g)))$ holds for every g in $(\mathcal{S}^\sharp)^2$. Let $g_1|g_2$ be an $\mathcal{S}$-normal path. By construction, we have $g_1 =^\times H(g_1 g_2)$. As ϕ_Δ is a functor, this implies $\phi_\Delta(g_1) \preccurlyeq \phi_\Delta(H(g_1 g_2))$ and $\phi_\Delta(H(g_1 g_2)) \preccurlyeq \phi_\Delta(g_1)$, hence $\phi_\Delta(g_1) =^\times \phi_\Delta(H(g_1 g_2))$. As we have $\phi_\Delta(H(g_1 g_2)) =^\times H(\phi_\Delta(g_1 g_2))$ by assumption, we deduce $\phi_\Delta(g_1) =^\times H(\phi_\Delta(g_1 g_2))$. Hence $\phi_\Delta(g_1)$ is an $\mathcal{S}$-head for $\phi_\Delta(g_1 g_2)$. As we have $\phi_\Delta(g_1 g_2) = \phi_\Delta(g_1)\,\phi_\Delta(g_2)$, we deduce that $\phi_\Delta(g_1)|\phi_\Delta(g_2)$ is an $\mathcal{S}$-normal decomposition of $\phi_\Delta(g_1 g_2)$. Hence ϕ_Δ preserves normality.

(ii) Assume that ϕ_Δ preserves left-gcds on $\mathcal{S}^\sharp$. Let $g_1|g_2$ be an $\mathcal{S}$-normal path. Let x be the source of g_2. By Proposition V.1.53, $\partial_\Delta(g_1)$ and g_2 are left-coprime, that

is, 1_x is a left-gcd of $\partial_\Delta(g_1)$ and g_2. If the condition of the statement is satisfied, it follows that $\phi_\Delta(1_x)$, which is $1_{\phi_\Delta(x)}$, is a left-gcd of $\phi_\Delta(\partial_\Delta(g_1))$ and $\phi_\Delta(g_2)$. By (V.1.30), we have $\phi_\Delta(\partial_\Delta(g_1)) = \partial_\Delta(\phi_\Delta(g_1))$. So $\partial_\Delta(\phi_\Delta(g_1))$ and $\phi_\Delta(g_2)$ are left-coprime, hence, by Proposition V.1.53, $\phi_\Delta(g_1)|\phi_\Delta(g_2)$ is $\mathcal{S}$-normal.

Exercice 62 (**normal decomposition**).— *Assume that $\mathcal{C}$ is a left-cancellative category, Δ is a right-Garside map in $\mathcal{C}$ such that ϕ_Δ preserves normality, and f, g are elements of $\mathcal{C}$ such that fg is defined and $f \preccurlyeq \Delta^{[m]}(\text{-})$ holds, say $ff' = \Delta^{[m]}(\text{-})$ with $m \geqslant 1$. Show that f' and g admit a left-gcd and that, if h is such a left-gcd, then concatenating a $\mathcal{D}iv(\Delta)$-normal decomposition of fh and a $\mathcal{D}iv(\Delta)$-normal decomposition of $h^{-1}g$ yields a $\mathcal{D}iv(\Delta)$-normal decomposition of fg. [Hint: First show that concatenating fh and a $\mathcal{D}iv(\Delta^{[m]})$-normal decomposition of $h^{-1}g$ yields a $\mathcal{D}iv(\Delta^{[m]})$-normal decomposition of fg and apply Exercise 42 in Chapter IV.]*

Solution. By Proposition V.1.58, $\Delta^{[m]}$ is a right-Garside map in $\mathcal{C}$ and, by definition, f' right-divides an element $\Delta^{[m]}(\text{-})$ so, by Lemma V.2.36, f' and g admit a left-gcd. Assume that h is a left-gcd of f' and g. Since $\Delta^{[m]}$ is a right-Garside map, gh and $\Delta^{[m]}(\text{-})$, that is, ff', admit a left-gcd, which (by Exercise 8(i) in Chapter II) is of the form fh' with h' a left-gcd of f' and g. By uniqueness, we have $h =^\times h'$, whence $fh =^\times fh'$. So fh is a left-gcd of fg and $\Delta^{[m]}(\text{-})$, hence it is a $\mathcal{D}iv(\Delta^{[m]})$-head of fg. Hence concatenating fh and a $\mathcal{D}iv(\Delta^{[m]})$-normal decomposition of $h^{-1}g$ yields a $\mathcal{D}iv(\Delta^{[m]})$-normal decomposition of fg. Then Exercise 42 gives the result.

Exercice 64 (**iterated duality**).— *Assume thatΔ is a Garside map in a cancellative category $\mathcal{C}$.* (i) *Show that $\partial^{[m']}(g) = \partial^{[m]}(g)\,\Delta^{[m'-m]}(\phi_\Delta^m(x))$ holds for $m' \geqslant m$ and g in $\mathrm{Div}(\Delta^{[m]}(x))$.* (ii) *Show that $\widetilde{\partial}_{[m']}(g) = \widetilde{\Delta}_{[m'-m]}(\phi_\Delta^{-m}(y))\,\widetilde{\partial}_{[m]}(g)$ holds for $m' \geqslant m$ and g in $\widetilde{\mathrm{Div}}(\widetilde{\Delta}_{[m]}(y))$.*

Solution. (i) By definition, we have

$$g\,\partial^{[m]}(g)\,\Delta^{[m'-m]}(\phi_\Delta^m(x)) = \Delta^{[m]}(x)\,\Delta^{[m'-m]}(\phi_\Delta^m(x)) = \Delta^{[m']}(x) = g\,\partial^{[m']}(g),$$

whence the result by left-cancelling g. The computation is symmetric for (ii).

Chapter VI: Germs

Skipped proofs

(none)

Solutions to selected exercises

Exercice 65 (**not embedding**).— *Let $\mathcal{S}$ consist of fourteen elements $1, \mathtt{a}, \dots, \mathtt{n}$, all with the same source and target, and $\bullet$ be defined by $1 \bullet x = x \bullet 1 = x$ for each x, plus $\mathtt{a}\bullet\mathtt{b} = \mathtt{f}$, $\mathtt{f}\bullet\mathtt{c} = \mathtt{g}$, $\mathtt{d}\bullet\mathtt{e} = \mathtt{h}$, $\mathtt{g}\bullet\mathtt{h} = \mathtt{i}$, $\mathtt{c}\bullet\mathtt{d} = \mathtt{j}$, $\mathtt{b}\bullet\mathtt{j} = \mathtt{k}$, $\mathtt{k}\bullet\mathtt{e} = \mathtt{m}$, and $\mathtt{a}\bullet\mathtt{m} = \mathtt{n}$.* (i) *Shows that $\underline{\mathcal{S}}$ is a germ.* (ii) *Show that, in $\mathcal{S}$, we have $((\mathtt{a}\bullet\mathtt{b})\bullet\mathtt{c})\bullet(\mathtt{d}\bullet\mathtt{e}) = \mathtt{i} \neq \mathtt{n} = \mathtt{a}\bullet((\mathtt{b}\bullet(\mathtt{c}\bullet\mathtt{d}))\bullet\mathtt{e})$, whereas, in $\mathcal{M}on(\underline{\mathcal{S}})$, we have $\iota\mathtt{i} = \iota\mathtt{n}$. Conclude.*

Solution. (i) The only nontrivial triples eligible for (VI.1.6) are $(\mathtt{a},\mathtt{b},\mathtt{j})$, $(\mathtt{f},\mathtt{c},\mathtt{d})$, and $(\mathtt{c},\mathtt{d},\mathtt{e})$, for which (VI.1.6) is actually true. (ii) The equalities in $\underline{\mathcal{S}}$ follow from the multiplication table of $\underline{\mathcal{S}}$. On the other hand, in $\mathcal{M}on(\underline{\mathcal{S}})$, we find $\iota\mathtt{i} = ((\iota\mathtt{a}\,\iota\mathtt{b})\,\iota\mathtt{c})\,(\iota\mathtt{d}\,\iota\mathtt{e}) = \iota\mathtt{a}\,((\iota\mathtt{b}\,(\iota\mathtt{c}\,\iota\mathtt{d}))\,\iota\mathtt{e}) = \iota\mathtt{n}$ applying associativity. Thus ι is not injective, and $\mathcal{S}^\sharp$ does not embed in $\mathcal{M}on(\underline{\mathcal{S}})$.

Exercice 66 (**multiplying by invertible elements**).— (i) *Show that, if $\underline{\mathcal{S}}$ is a left-associative germ, then $\mathcal{S}$ is closed under left-multiplication by invertible elements in $\mathcal{C}at(\underline{\mathcal{S}})$.* (ii) *Show that, if $\underline{\mathcal{S}}$ is an associative germ, $s\bullet t$ is defined, and $t' =^{\times}_{\mathcal{S}} t$ holds, then $s\bullet t'$ is defined as well.*

Solution. (i) Assume that ϵ admits a left-inverse ϵ'—as nothing a priori forces the category $\mathcal{C}at(\underline{\mathcal{S}})$ to be left-cancellative, we have to distinguish between left- and right-inverse—and that ϵg is defined for some g lying in $\mathcal{S}$. Then we have $g = \epsilon'\epsilon g$, so ϵg is a right-divisor of an element of $\mathcal{S}$, hence is an element of $\mathcal{S}$ as the latter is closed under right-divisor. Applying this to the case when g is an identity-element shows that the family of all left-invertible elements is included in $\mathcal{S}$.

Exercice 67 (**atoms**).— (i) *Show that, if $\underline{\mathcal{S}}$ is a left-associative germ, the atoms of $\mathcal{C}at(\underline{\mathcal{S}})$ are the elements of the form $t\epsilon$ with t an atom of $\underline{\mathcal{S}}$ and ϵ an invertible element of $\underline{\mathcal{S}}$.*

(ii) *Let $\underline{\mathcal{S}}$ be the germ whose table is shown on the right. Show that the monoid $\mathcal{M}on(\underline{\mathcal{S}})$ admits the presentation $\langle \mathtt{a},\mathtt{e} \mid \mathtt{ea} = \mathtt{a}, \mathtt{e}^2 = 1\rangle^+$ (see Exercise 48) and that $\mathtt{a}$ is the only atom of $\underline{\mathcal{S}}$, whereas the atoms of $\mathcal{M}on(\underline{\mathcal{S}})$ are $\mathtt{a}$ and $\mathtt{ae}$.* (iii) *Show that $\underline{\mathcal{S}}$ is a Garside germ.*

$\bullet$	1	a	e
1	1	a	e
a	a		
e	e	a	1

Solution. (i) Assume $s = t\epsilon$ with t an atom of $\underline{\mathcal{S}}$ and ϵ an invertible element in $\mathcal{S}$. Let $s_1|\cdots|s_p$ be a decomposition of s in $\mathcal{C}at(\underline{\mathcal{S}})$. As $\mathcal{S}$ generates $\mathcal{C}at(\underline{\mathcal{S}})$, each element s_i can be expressed as a product of elements of $\mathcal{S}$, leading to a new decomposition $t_1|\cdots|t_q$ of s in terms of elements of $\mathcal{S}$, whence $t = t_1\cdots t_q\epsilon^{-1}$ in $\mathcal{C}at(\underline{\mathcal{S}})$ and, as $\underline{\mathcal{S}}$ is left-associative, $t = \Pi(t_1|\cdots|t_q|\epsilon^{-1})$ in $\underline{\mathcal{S}}$, where Π is the partial map of (VI.1.15). As t is an atom of $\underline{\mathcal{S}}$, at most one of the entries t_j is non-invertible in $\underline{\mathcal{S}}$. Hence at most one of the entries s_i is non-invertible in $\mathcal{C}at(\underline{\mathcal{S}})$, and s is an atom in $\mathcal{C}at(\underline{\mathcal{S}})$.

Conversely, assume that s is an atom in $\mathcal{C}at(\underline{\mathcal{S}})$. As $\mathcal{S}$ generates $\mathcal{C}at(\underline{\mathcal{S}})$, there exists a decomposition $s_1|\cdots|s_p$ of s into elements of $\mathcal{S}$. As s is an atom in $\mathcal{C}at(\underline{\mathcal{S}})$, at most one entry is not invertible. If every entry is invertible, then s is invertible, contradicting the assumption. So exactly one entry, say s_i, is not invertible. Let $t = s_1\cdots s_{i-1}s_i$ and $\epsilon = s_{i+1}\cdots s_p$. As $s_1,\dots,s_{i-1}$ are invertible, t is a right-divisor of s, hence it belongs to $\mathcal{S}$. Moreover, t is an atom of $\mathcal{S}$, since a decomposition of t with more than one non-invertible entry in $\mathcal{S}$ would provide a similar decomposition in $\mathcal{C}at(\underline{\mathcal{S}})$, contradicting the assumption. On the other hand, ϵ is invertible. So s, which is $t\epsilon$, has the expected form.

Exercice 68 (**families $\mathcal{I}_{\underline{\mathcal{S}}}$ and $\mathcal{J}_{\underline{\mathcal{S}}}$**).— *Assume that $\underline{\mathcal{S}}$ is a left-associative germ.* (i) *Show that a path $s_1|s_2$ of $\mathcal{S}^{[2]}$ is $\mathcal{S}$-normal if and only if all elements of $\mathcal{J}_{\underline{\mathcal{S}}}(s_1,s_2)$ are invertible.* (ii) *Assuming in addition that $\underline{\mathcal{S}}$ is left-cancellative, show that, for $s_1|s_2$ in $\mathcal{S}^{[2]}$, the family $\mathcal{I}_{\underline{\mathcal{S}}}(s_1,s_2)$ admits common right-multiples if and only if $\mathcal{J}_{\underline{\mathcal{S}}}(s_1,s_2)$ does.*

Solution. (ii) Assume that $\mathfrak{I}_{\underline{\mathcal{S}}}(g_1, g_2)$ admits common right-multiples, and let h, h' belong to $\mathfrak{J}_{\underline{\mathcal{S}}}(g_1, g_2)$. Then $g_1 \bullet h$ and $g_2 \bullet h'$ are defined and belong to $\mathfrak{I}_{\underline{\mathcal{S}}}(g_1, g_2)$, hence, by assumption, $g_1 \bullet h$ and $g_1 \bullet h'$ admit a common right-multiple, say g''. Then we have $g_1 \preccurlyeq_{\mathcal{S}} g_1 \bullet h \preccurlyeq_{\mathcal{S}} g$, whence $g_1 \preccurlyeq_{\mathcal{S}} g''$. So there exists h'' in $\mathcal{S}$ satisfying $g'' = g_1 \bullet h''$. By Lemma VI.1.19, $g_1 \bullet h \preccurlyeq_{\mathcal{S}} g_1 \bullet h''$ implies $h \preccurlyeq_{\mathcal{S}} h''$, and, similarly, we find $h' \preccurlyeq_{\mathcal{S}} h''$. So h'' is a common right-multiple of h and h' in $\mathcal{S}$. Moreover the assumption that $g_1 \bullet h''$ belongs to $\mathfrak{I}_{\underline{\mathcal{S}}}(g_1, g_2)$ implies that h'' belongs to $\mathfrak{J}_{\underline{\mathcal{S}}}(g_1, g_2)$. So $\mathfrak{J}_{\underline{\mathcal{S}}}(g_1, g_2)$ admits common right-multiples. The converse implication is similar, actually simpler as no cancellation is needed.

Exercice 69 (**positive generators**).— *Assume that Σ is a family of positive generators in a group $\mathcal{G}$ and Σ is closed under inverse, that is, $g \in \Sigma$ implies $g^{-1} \in \Sigma$.* (i) *Show that $\|g\|_\Sigma = \|g^{-1}\|_\Sigma$ holds for every g in $\mathcal{G}$.* (ii) *Show that $f^{-1} \leqslant_\Sigma g^{-1}$ is equivalent to $f \widetilde{\leqslant}_\Sigma g$.*

Solution. (i) An S-word w is a minimal length expression for an element g if and only if w^{-1}, which is also an S-word, is a minimal length expression of g^{-1}. (ii) By definition, $f^{-1} \leqslant_\Sigma g^{-1}$ is equivalent to $\|f^{-1}\|_\Sigma + \|(f^{-1})^{-1} g^{-1}\|_\Sigma = \|g^{-1}\|_\Sigma$, hence, by (i), to $\|f\|_\Sigma + \|f g^{-1}\|_\Sigma = \|g\|_\Sigma$ and to $\|f\|_\Sigma + \|g f^{-1}\|_\Sigma = \|g\|_\Sigma$. The latter is $f \widetilde{\leqslant}_\Sigma g$.

Exercice 70 (**minimal upper bound**).— *For $\leqslant$ a partial ordering on a family $\mathcal{S}'$ and f, g, h in $\mathcal{S}'$, say that h is a* minimal upper bound*, or* mub*, for f and g, if $f \leqslant h$ and $g \leqslant h$ holds, but there exists no h' with $h' < h$ satisfying $f \leqslant h'$ and $g \leqslant h'$. Assume that $\mathcal{G}$ is a groupoid, Σ positively generates $\mathcal{G}$, and $\mathcal{H}$ is a subfamily of $\mathcal{G}$ that is closed under Σ-suffix. Show that $\mathcal{H}^\Sigma$ is a Garside germ if and only if, for all f, g, g', g'' in $\mathcal{H}$ such that $f \bullet g$ and $f \bullet g'$ are defined and g'' is a $\leqslant_\Sigma$-mub of g and g', the product $f \bullet g''$ is defined.*

Solution. By Lemmas VI.2.60 and VI.2.62, the germ $\mathcal{H}^\Sigma$ is left-associative, cancellative, and Noetherian. Hence, by Proposition VI.2.44, $\mathcal{H}^\Sigma$ is a Garside germ if and only if it satisfies (VI.2.43). Now, by Lemma VI.2.62, for g, h in $\mathcal{H}$, the relation $g \preccurlyeq_{\mathcal{H}^\Sigma} h$ is equivalent to $g \leqslant_\Sigma h$ and, therefore, g'' is an mcm of g and g' in $\mathcal{H}^\Sigma$ if and only it it is a $\leqslant_\Sigma$-mub of g and g'. So the condition is a direct reformulation of(VI.2.43).

Chapter VII: Subcategories

Skipped proofs

Lemma VII.1.3.— *If $\mathcal{C}_1$ is a subcategory of a left-cancellative category $\mathcal{C}$, we have*

$$\mathcal{C}_1^\times \subseteq \mathcal{C}^\times \cap \mathcal{C}_1, \tag{VII.1.4}$$

with equality if and only if $\mathcal{C}_1$ is closed under inverse in $\mathcal{C}$. For every subfamily $\mathcal{S}$ of $\mathcal{C}$, putting $\mathcal{S}_1 = \mathcal{S} \cap \mathcal{C}_1$ and $\mathcal{S}_1^\sharp = \mathcal{S}_1 \mathcal{C}_1^\times \cup \mathcal{C}_1^\times$, we have

$$\mathcal{S}_1^\sharp \subseteq \mathcal{S}^\sharp \cap \mathcal{C}_1. \tag{VII.1.5}$$

If $\mathcal{C}$ has no nontrivial invertible element, then so does $\mathcal{C}_1$, and (VII.1.4)–(VII.1.5) are equalities.

Proof. If $\epsilon\epsilon' = 1_x$ holds in $\mathcal{C}_1$, it holds in $\mathcal{C}$ as well, so (VII.1.4) is clear. For (VII.1.4) to be an equality means that every invertible element lying in $\mathcal{C}_1$ belongs to $\mathcal{C}_1^\times$, that is, has an inverse that lies in $\mathcal{C}_1$: this means that $\mathcal{C}_1$ is closed under inverse in $\mathcal{C}$.

Next, assume that $\mathcal{S}$ is included in $\mathcal{C}$, and put $\mathcal{S}_1 = \mathcal{S} \cap \mathcal{C}_1$ and $\mathcal{S}_1^\sharp = \mathcal{S}_1\mathcal{C}_1^\times \cap \mathcal{C}_1^\times$. Then $\mathcal{C}_1^\times$ is included in $\mathcal{C}^\times$ and in $\mathcal{C}_1$, so we deduce $\mathcal{S}_1^\sharp \subseteq (\mathcal{S}\mathcal{C}^\times \cup \mathcal{C}^\times) \cap \mathcal{C}_1 = \mathcal{S}^\sharp \cap \mathcal{C}_1$.

On the other hand, assume $\mathcal{C}^\times = \mathbf{1}_\mathcal{C}$. As an identity-element is its own inverse, it is invertible in every subcategory that contains it, and we obtain $\mathcal{C}_1^\times = \mathbf{1}_\mathcal{C} \cap \mathcal{C}_1 = \mathcal{C}^\times \cap \mathcal{C}_1$ and, for every $\mathcal{S} \subseteq \mathcal{C}$, as $\mathcal{S}^\sharp$ is then $\mathcal{S} \cup \mathbf{1}_\mathcal{C}$, we obtain

$$\mathcal{S}^\sharp \cap \mathcal{C}_1 = (\mathcal{S} \cap \mathcal{C}_1) \cup (\mathbf{1}_\mathcal{C} \cap \mathcal{C}_1) = \mathcal{S}_1 \cup \mathcal{C}_1^\times = \mathcal{S}_1\mathcal{C}_1^\times \cup \mathcal{C}_1^\times = \mathcal{S}_1^\sharp. \qquad \square$$

Lemma VII.1.16.— *Every subcategory that is closed under left- or under right-divisor in a left-cancellative category $\mathcal{C}$ is closed under $=^\times$.*

Proof. Assume that $\mathcal{C}_1$ is a subcategory of $\mathcal{C}$, and we have $g \in \mathcal{C}_1$ and $g' =^\times g$, say $g' = g\epsilon$ with ϵ invertible. If $\mathcal{C}_1$ is closed under left-divisor, we can write $g = g'\epsilon^{-1}$, and g' is a left-divisor of g, hence it belongs to $\mathcal{C}_1$.

On the other hand, assume that $\mathcal{C}_1$ is closed under right-divisor. Let y be the target of g, and y' be that of g'. The assumption that g lies in $\mathcal{C}_1$ implies that y lies in $\mathcal{O}bj(\mathcal{C}_1)$, hence 1_y lies in $\mathcal{C}_1$. Now ϵ^{-1} is a right-divisor of g, hence it lies in $\mathcal{C}_1$, its source y' lies in $\mathcal{O}bj(\mathcal{C}_1)$ and $1_{y'}$ lies in $\mathcal{C}_1$. Now, we have $1_{y'} = \epsilon^{-1}\epsilon$, so ϵ, a right-divisor of 1_z, must lie in $\mathcal{C}_1$. Hence g', that is, $g\epsilon$, lies in $\mathcal{C}_1$. $\square$

Lemma VII.1.17.— *If $\mathcal{C}, \mathcal{C}'$ are left-cancellative categories, $\phi : \mathcal{C} \to \mathcal{C}'$ is a functor, and $\mathcal{C}_1'$ is a subcategory of $\mathcal{C}'$ that is closed under left-divisor (resp. right-divisor), then so is $\phi^{-1}(\mathcal{C}_1')$.*

Proof. Assume that $\mathcal{C}_1'$ is closed under left-divisor. Put $\mathcal{C}_1 = \phi^{-1}(\mathcal{C}_1')$, and assume $f \preccurlyeq g \in \mathcal{C}_1$. By definition, there exists g' satisfying $fg' = g$, which implies $\phi(f)\phi(g') = \phi(g)$, whence $\phi(f) \preccurlyeq \phi(g)$. By assumption, $\phi(g)$ lies in $\mathcal{C}_1'$, hence so does $\phi(f)$ as $\mathcal{C}_1'$ is closed under left-divisor. Hence $\phi(f)$ belongs to $\mathcal{C}_1'$, and f lies in $\mathcal{C}_1$. So $\mathcal{C}_1$ is closed under left-divisor. The argument when $\mathcal{C}_1'$ is closed under right-divisor is symmetric. $\square$

Proposition VII.2.16 (**recognizing compatible**).— *If $\mathcal{S}$ is a Garside family in a left-cancellative category $\mathcal{C}$ and $\mathcal{C}_1$ is a subcategory of $\mathcal{C}$ that is closed under right-quotient in $\mathcal{C}$, then $\mathcal{C}_1$ is compatible with $\mathcal{S}$ if and only if*

(VII.2.17) *The family $\mathcal{S}_1^\sharp$ generates $\mathcal{C}_1$, where we put $\mathcal{S}_1 = \mathcal{S} \cap \mathcal{C}_1$ and $\mathcal{S}_1^\sharp = \mathcal{S}_1\mathcal{C}_1^\times \cup \mathcal{C}_1^\times$,*

(VII.2.18) *Every element of $(\mathcal{S}_1^\sharp)^2$ admits an $\mathcal{S}$-normal decomposition with entries in $\mathcal{S}_1^\sharp$.*

Proof. If $\mathcal{C}_1$ is compatible with $\mathcal{S}$, then, by Proposition VII.2.14, (VII.2.15) holds. This implies in particular that every element of $\mathcal{C}_1$ admits a decomposition with entries in $\mathcal{S}_1^\sharp$, hence that $\mathcal{S}_1^\sharp$ generates $\mathcal{C}_1$, which is (VII.2.17). On the other hand, applying (VII.2.15) to an element of $(\mathcal{S}_1^\sharp)^2$ gives (VII.2.18).

Conversely, assume that (VII.2.17) and (VII.2.18) are satisfied. As $\mathcal{C}_1$ is closed under right-quotient, it is closed under inverse, $\mathcal{S}_1^\sharp \cap \mathcal{C}^\times = \mathcal{C}_1^\times$ holds, hence so does

$\mathcal{S}_1^\sharp(\mathcal{S}_1^\sharp \cap \mathcal{C}^\times) \subseteq \mathcal{S}_1^\sharp$. Then Lemma VII.2.19 is valid for $\mathcal{S}_1^\sharp$, so every element of the subcategory of $\mathcal{C}$ generated by $\mathcal{S}_1^\sharp$, hence of $\mathcal{C}_1$, admits an $\mathcal{S}$-normal decomposition whose entries lie in $\mathcal{S}_1^\sharp$. In this case, (VII.2.15) is satisfied, and, by Proposition VII.2.14, $\mathcal{C}_1$ is compatible with $\mathcal{S}$. □

Lemma VII.2.19.— *Assume that $\mathcal{S}$ is a Garside family in a left-cancellative category $\mathcal{C}$ and $\mathcal{S}'$ is a subfamily of $\mathcal{S}^\sharp$ such that $\mathcal{S}'(\mathcal{S}' \cap \mathcal{C}^\times) \subseteq \mathcal{S}'$ holds and every element of $\mathcal{S}'^2$ admits a $\mathcal{S}$-normal decomposition with entries in $\mathcal{S}'$. Then every element in the subcategory of $\mathcal{C}$ generated by $\mathcal{S}'$ admits a $\mathcal{S}$-normal decomposition with entries in $\mathcal{S}'$.*

Proof. First we claim that every element g of $\mathcal{S}'^2$ admits a length two $\mathcal{S}$-normal decomposition with entries in $\mathcal{S}'$: indeed, as $\mathcal{S}'$ is included in $\mathcal{S}^\sharp$, the $\mathcal{S}$-length of g is at most two, so the possible entries at position 3 and beyond in an $\mathcal{S}$-normal decomposition of g must be invertible, and the assumption $\mathcal{S}'(\mathcal{S}' \cap \mathcal{C}^\times) \subseteq \mathcal{S}'$ implies that the latter can be incorporated in the second entry, yielding an $\mathcal{S}$-normal decomposition of length two.

Then the argument is exactly the same as for Proposition III.1.49 (left multiplication), except that, here, we use two different reference families, namely $\mathcal{S}'$ for the entries of the decomposition and $\mathcal{S}$ for the greedy property. The point is to prove that, for every p, every element of $\mathcal{S}'^p$ admits an $\mathcal{S}$-normal decomposition of length p with entries in $\mathcal{S}'$. We use induction on $p \geqslant 1$. For $p = 1$, the result is obvious as $\mathcal{S}'$ is included in $\mathcal{S}^\sharp$, and, for $p = 2$, the result is the assumption. Let g belong to $\mathcal{S}'^p$ with $p \geqslant 3$. Write $g = sg'$ with $s \in \mathcal{S}'$ and $g' \in \mathcal{S}'^{p-1}$. By induction hypothesis, g' admits an $\mathcal{S}$-normal decomposition $s'_1|\cdots|s'_{p-1}$ with entries in $\mathcal{S}'$. Starting with $s_0 = s$, and applying the assumption $p-1$ times, we find an $\mathcal{S}$-normal decomposition $g_i|s_i$ of $s_{i-1}g'_i$ with entries in $\mathcal{S}'$. By the first domino rule (Proposition III.1.45), $s_1|\cdots|s_{p-1}|s_p$ is an $\mathcal{S}$-normal decomposition of g with entries in $\mathcal{S}'$. □

Lemma VII.4.7.— *If $\mathcal{S}$ is any subfamily of a left-cancellative category $\mathcal{C}$, the identity-functor on $\mathcal{C}$ is correct for inverses (resp. right-comultiples, resp. right-complements, resp. right-diamonds) on $\mathcal{S}$ if and only if $\mathcal{S}$ is closed under inverse (resp. right-comultiple, resp. right-complement, resp. right-diamond) in $\mathcal{C}$.*

Proof. The identity-functor on $\mathcal{C}$ is correct for inverses on $\mathcal{S}_1$ if and only if, for every s in $\mathcal{S}_1$ that is invertible, s^{-1} belongs to $\mathcal{S}_1$. By definition, this means that $\mathcal{S}_1$ is closed under inverse in $\mathcal{C}$.

Next, the identity-functor on $\mathcal{C}$ is correct for right-comultiples on $\mathcal{S}_1$ if and only if, when s, t lie in $\mathcal{S}_1$ and $sg = tf$ holds in $\mathcal{C}$, there exist s', t', and h satisfying $st' = ts'$, $f = s'h$, and $g = t'h$, plus $st' \in \mathcal{S}_1$. By definition, this means that $\mathcal{S}_1$ is closed under right-comultiple in $\mathcal{C}$. The result is similar with right-complements and right-diamonds. □

Solutions to selected exercises

Exercice 72 (**$=^\times$-closed subcategory**).— (i) *Show that a subcategory $\mathcal{C}_1$ of a left-cancellative category $\mathcal{C}$ is $=^\times$-closed if and only if, for each x in $\mathcal{O}bj(\mathcal{C}_1)$, the*

families $\mathcal{C}^\times(x,\text{-})$ and $\mathcal{C}^\times(\text{-},x)$ are included in $\mathcal{C}_1$. (ii) *Deduce that $\mathcal{C}_1$ is $=^\times$-closed if and only if $\mathcal{C}_1^\times$ is a union of connected components of $\mathcal{C}^\times$.*

Solution. Assume that $\mathcal{C}_1$ is $=^\times$-closed and x lies in $\mathcal{O}bj(\mathcal{C}_1)$. Then 1_x belongs to $\mathcal{C}_1$. If ϵ belongs to $\mathcal{C}^\times(x,\text{-})$, we have $\epsilon =^\times 1_x$, whence $\epsilon \in \mathcal{C}_1$. On the other hand, if ϵ belongs to $\mathcal{C}^\times(y,x)$, then ϵ^{-1} belongs to $\mathcal{C}^\times(x,y)$. By the above result, ϵ^{-1} belongs to $\mathcal{C}_1$. It follows that y, the target of ϵ^{-1}, lies in $\mathcal{O}bj(\mathcal{C}_1)$, and, therefore, ϵ belongs to $\mathcal{C}_1$. Conversely, assume that $\mathcal{C}^\times(x,\text{-})$ is included in $\mathcal{C}_1$ for every x in $\mathcal{O}bj(\mathcal{C}_1)$, and let g belong to $\mathcal{C}_1$ and $g' =^\times g$ hold, say $g' = g\epsilon$ with ϵ in $\mathcal{C}^\times$. Let x be the target of g. Then x belongs to $\mathcal{O}bj(\mathcal{C}_1)$, hence ϵ belongs to $\mathcal{C}_1$, so $g \in \mathcal{C}_1$ implies $g'g\epsilon \in \mathcal{C}_1$.

Exercice 73 (**greedy paths**).— *Assume that $\mathcal{C}$ is a cancellative category, $\mathcal{S}$ is included in $\mathcal{C}$, and $\mathcal{C}_1$ is a subcategory of $\mathcal{C}$ that is closed under left-quotient. Put $\mathcal{S}_1 = \mathcal{S} \cap \mathcal{C}_1$. Show that every $\mathcal{C}_1$-path that is $\mathcal{S}$-greedy in $\mathcal{C}$ is $\mathcal{S}_1$-greedy in $\mathcal{C}_1$.*

Solution. With the notation of the proof of Lemma VII.2.1, we have also $f' = f''g_2$ with f' and g_2 in $\mathcal{C}_1$. This implies $f'' \in \mathcal{S}_1$ as $\mathcal{C}$ is right-cancellative and $\mathcal{C}_1$ is closed under left-quotient in $\mathcal{C}$.

Exercice 74 (**compatibility with $\mathcal{C}$**).— *Assume that $\mathcal{C}$ is a left-cancellative category. Show that every subcategory of $\mathcal{C}$ that is closed under inverse is compatible with $\mathcal{C}$ viewed as a Garside family in itself.*

Solution. Let $\mathcal{C}_1$ be a subcategory of $\mathcal{C}$ that is closed under inverse. Then $\mathcal{C} \cap \mathcal{C}_1 = \mathcal{C}_1$ is a Garside family in $\mathcal{C}_1$. A $\mathcal{C}_1$-path $g_1|\cdots|g_q$ is $\mathcal{C}$-normal if and only if $g_2, ..., g_p$ belong to $\mathcal{C}^\times$, hence if and only if $g_2, ..., g_p$ belong to $\mathcal{C}_1^\times$, hence if and only if $g_1|\cdots|g_q$ is $\mathcal{C}_1$-normal.

Exercice 75 (**not compatible**).— *Let M be the free Abelian monoid generated by $\mathtt{a}$ and $\mathtt{b}$, and let N be the submonoid generated by $\mathtt{a}$ and $\mathtt{ab}$.* (i) *Show that N is not closed under right-quotient in M.* (ii) *Let $S = \{1, \mathtt{a}, \mathtt{b}, \mathtt{ab}\}$. Show that N is not compatible with S.* (iii) *Let $S' = \{\mathtt{a}^p\mathtt{b}^\epsilon \mid p \geqslant 0, i \in \{0,1\}\}$. Show that N not compatible with S'.*

Solution. (i) The elements $\mathtt{ab}$ and $\mathtt{a}$ lie in N, but the right-quotient $\mathtt{b}$ does not.

(ii) The family $S \cap N$ is not a Garside family in N, because $\mathtt{a}$ and $\mathtt{ab}$, which belong to $S \cap N$, have no common right-multiple belonging to $S \cap N$.

(iii) The family $S' \cap N$ is a Garside family in M, but the S'-normal decomposition of $\mathtt{a}^2\mathtt{b}^2$ is $\mathtt{a}^2\mathtt{b}|\mathtt{b}$, whereas the $(S' \cap N)$-normal decomposition of $\mathtt{a}^2\mathtt{b}^2$ in N is $\mathtt{ab}|\mathtt{ab}$.

Exercice 76 (**not closed under right-quotient**).— (i) *Show that every submonoid $m\mathbb{N}$ of the additive monoid $\mathbb{N}$ is closed under right-quotient, but that $2\mathbb{N}+3\mathbb{N}$ of $\mathbb{N}$ is not.* (ii) *Let M be the monoid $\mathbb{N} \rtimes (\mathbb{Z}/2\mathbb{Z})^2$, where the generator $\mathtt{a}$ of $\mathbb{N}$ acts on the generators $\mathtt{e}, \mathtt{f}$ of $(\mathbb{Z}/2\mathbb{Z})^2$ by $\mathtt{ae} = \mathtt{fa}$ and $\mathtt{af} = \mathtt{ea}$, and let N be the submonoid of M generated by $\mathtt{a}$ and $\mathtt{e}$. Show that M is left-cancellative, and its elements admit a unique expression of the form $\mathtt{a}^p\mathtt{e}^i\mathtt{f}^j$ with $p \geqslant 0$ and $i, j \in \{0,1\}$, and that N is $M \setminus \{\mathtt{f}, \mathtt{ef}\}$.* (iii) *Show that N is not closed under right-quotient in M.* (iv) *Let $S = \{\mathtt{a}\}$. Show that S is a Garside family in M and determine $S^\sharp$. Show that N is compatible with S. [Hint: Show that $S \cap N$, which is S, is not a Garside family in N.]* (v) *Shows that $S^\sharp \cap N$ is a Garside family in N and N is compatible with $S^\sharp$.*

Solution. (i) If f and $f+g'$ are multiples of m, then g' is a multiple of m as well. On the other hand, $2\mathbb{N}+3\mathbb{N}$ contains 2 and 3, but does not contain 1 although $3=2+1$ holds.

(iii) The set $M^\times$ consists of $1, \mathtt{e}, \mathtt{f}, \mathtt{ef}$, whereas $N^\times$ consists of 1 and $\mathtt{e}$, so N is closed under inverse in M. On the other hand, as $\mathtt{a}$ and $\mathtt{af}$ belong to N but $\mathtt{f}$ does not, N is not closed under right-quotient in M.

(iv) We have $S^\sharp = \{\mathtt{a}^p\mathtt{e}^i\mathtt{f}^j \mid p,i,j \leqslant 1\}$ (eight elements). Then $\mathtt{af}$, which is $\mathtt{ea}$, belongs to $N^\times S$ but not to $SN^\times \cup N^\times$. So $S\cap N$, which is S, is not a Garside family in N, and the submonoid N is not compatible with S.

(v) Consider $S^\sharp$, which is also a Garside family in M. Then $S^\sharp \cap N$ is $S^\sharp \setminus \{\mathtt{f}, \mathtt{ef}\}$ (six elements), and it is equal to $(S^\sharp \cap N)N^\times \cup N^\times$. Hence $S^\sharp \cap N$ generates N. Next, M and N are Noetherian, and both admit right-lcms. Now a direct inspection shows that $S^\sharp \cap N$ is (weakly) closed under right-lcm and right-divisor in N (here we consider only right-divisors that belong to N, so $\mathtt{f}$, which is a right-divisor of $\mathtt{af}$ in M, is excluded). Hence, by Corollary IV.2.29 (recognizing Garside, right-lcm case), $S^\sharp \cap N$ is a Garside family in N. Finally, a pair $\mathtt{a}^{p_1}\mathtt{e}^{i_1}\mathtt{f}^{j_1} | \mathtt{a}^{p_2}\mathtt{e}^{i_2}\mathtt{f}^{j_2}$ is S-normal in M if and only if we do not have $p_1=0$ and $p_2=1$ and the same condition characterizes $(S^\sharp\cap N)$-normal pairs in N. Hence N is compatible with $S^\sharp$.

Exercice 77 (**not closed under divisor**).— *Let $M = \langle \mathtt{a}, \mathtt{b} \mid \mathtt{ab} = \mathtt{ba}, \mathtt{a}^2 = \mathtt{b}^2\rangle^+$, and let N be the submonoid of M generated by $\mathtt{a}^2$ and $\mathtt{ab}$. Show that N is compatible with every Garside family S of M, but that M is not closed under left- and right-divisor.*

Solution. As seen in Example IV.2.34 (no proper Garside), the only Garside family in M is M itself.

Exercice 78 (**head implies closed**).— *Assume that $\mathcal{C}$ is a left-cancellative category, $\mathcal{S}$ is a subfamily of $\mathcal{C}$ that is closed under right-comultiple in $\mathcal{C}$, and $\mathcal{C}_1$ is a subcategory of $\mathcal{C}$. Put $\mathcal{S}_1 = \mathcal{S}\cap\mathcal{C}_1$.* (i) *Show that, if every element of $\mathcal{S}$ admits a $\mathcal{C}_1$-head that lies in $\mathcal{S}_1$, then $\mathcal{S}_1$ is closed under right-comultiple in $\mathcal{C}$.* (ii) *Show that, if, moreover, $\mathcal{S}_1$ is closed under right-quotient in $\mathcal{C}$, then $\mathcal{S}_1$ is closed under right-diamond in $\mathcal{C}$.*

Solution. (See Figure 9.) (i) Assume $sg = tf$ with $s,t \in \mathcal{S}_1$. As $\mathcal{S}$ is closed under right-comultiple, there exist s', t' satisfying $st' = ts' \preccurlyeq sg$ with $st' \in \mathcal{S}$. Let r be a $\mathcal{C}_1$-head of st' that lies in $\mathcal{S}_1$. As s lies in $\mathcal{C}_1$ and st' lies in $\mathcal{S}$, the relation $s \preccurlyeq st'$ implies $s \preccurlyeq r$, so we have $r = st''$ for some t'' in $\mathcal{C}_1$. Similarly, we have $r = ts_1$ for some s_1 in $\mathcal{C}_1$, and, therefore, r witnesses that $\mathcal{S}_1$ is closed under right-comultiple in $\mathcal{C}$. (ii) If, in addition, $\mathcal{S}_1$ is closed under right-quotient in $\mathcal{C}_1$, then s_1 and t_1 must lie in $\mathcal{S}_1$ since s,t, and r do, and $\mathcal{S}_1$ is closed under right-diamond in $\mathcal{C}$.

Exercice 79 (**head on generating family**).— *Assume that $\mathcal{C}$ is a left-cancellative category that is right-Noetherian, $\mathcal{C}_1$ is a subcategory of $\mathcal{C}$ that is closed under inverse, and $\mathcal{S}$ is a subfamily of $\mathcal{C}$ such that every element of $\mathcal{S}$ admits a $\mathcal{C}_1$-head that lies in $\mathcal{S}$. Assume moreover that $\mathcal{S}$ is closed under right-comultiple and that $\mathcal{S}\cap\mathcal{C}_1$ generates $\mathcal{C}_1$ and is closed under right-quotient in $\mathcal{C}$. Show that $\mathcal{C}_1$ is a head-subcategory of $\mathcal{C}$. [Hint: Apply Exercise 78.]*

Solution. Exercise 78 implies that $\mathcal{S}\cap\mathcal{C}_1$ is closed under right-diamond in $\mathcal{C}$. Then Proposition IV.1.15 (factorization grid) implies that the subcategory of $\mathcal{C}$



FIGURE 9. Solution to Exercise 79

generated by $\mathcal{S}\cap\mathcal{C}_1$, which is $\mathcal{C}_1$ by assumption, is closed under right-diamond in $\mathcal{C}$. By Lemma VII.1.8, it follows that $\mathcal{C}_1$ is closed under right-quotient in $\mathcal{C}$. Then, as $\mathcal{C}$ is right-Noetherian, Proposition VII.1.21 implies that $\mathcal{C}_1$ is a head-subcategory of $\mathcal{C}$.

Exercice 80 (**transitivity of compatibility**).— *Assume that $\mathcal{S}$ is a Garside family in a left-cancellative category $\mathcal{C}$, $\mathcal{C}_1$ is a subcategory of $\mathcal{C}$ that is compatible with $\mathcal{S}$, and $\mathcal{C}_2$ is a subcategory of $\mathcal{C}_1$ that is compatible with $\mathcal{S}_1 = \mathcal{S}\cap\mathcal{C}_1$. Show that $\mathcal{C}_2$ is compatible with $\mathcal{S}$.*

Solution. Put $\mathcal{S}_2 = \mathcal{S}_1\cap\mathcal{C}_2$. By assumption, we have $\mathcal{C}_2^\times = \mathcal{C}_1^\times\cap\mathcal{C}_2 = (\mathcal{C}^\times\cap\mathcal{C}_1)\cap\mathcal{C}_2 = \mathcal{C}^\times\cap\mathcal{C}_2$. Therefore, $\mathcal{C}_2$ in closed under inverse in $\mathcal{C}$. Next, as $\mathcal{C}_1$ is compatible with $\mathcal{S}$, the family $\mathcal{S}_1$ is a Garside family in $\mathcal{C}_1$. Now, as $\mathcal{C}_2$ is compatible with $\mathcal{S}_1$ in $\mathcal{C}_1$, the family $\mathcal{S}_2$, which is $\mathcal{S}_1\cap\mathcal{C}_2$, is a Garside family in $\mathcal{C}_2$. Finally, as $\mathcal{S}_2 = \mathcal{S}_1\cap\mathcal{C}_2$ holds, an $\mathcal{C}_2$ path is $\mathcal{S}_2$-normal in $\mathcal{C}_2$ if and only if it is $\mathcal{S}_1$-normal in $\mathcal{C}_1$, hence if and only if it is $\mathcal{S}$-normal in $\mathcal{C}$. Hence $\mathcal{C}_2$ is compatible with $\mathcal{S}$.

Exercice 81 (**transitivity of head-subcategory**).— *Assume that $\mathcal{C}$ is a left-cancellative category, $\mathcal{C}_1$ is a head-subcategory of $\mathcal{C}$, and $\mathcal{C}_2$ is a subcategory of $\mathcal{C}_1$. Show that $\mathcal{C}_2$ is a head-subcategory of $\mathcal{C}$ if and only if it is a head-subcategory of $\mathcal{C}_1$.*

Solution. Assume that $\mathcal{C}_2$ is a head-subcategory of $\mathcal{C}$. If ϵ belongs to $\mathcal{C}_1^\times\cap\mathcal{C}_2$, then it belongs to $\mathcal{C}^\times\cap\mathcal{C}_2$, hence ϵ^{-1} belongs to $\mathcal{C}_2$, so $\mathcal{C}_2$ is closed under inverse in $\mathcal{C}_1$. Moreover, every element of $\mathcal{C}_1$ admits a $\mathcal{C}_2$-head since every element of $\mathcal{C}$ does. Conversely assume that $\mathcal{C}_2$ is a head-subcategory of $\mathcal{C}_1$. If ϵ belongs to $\mathcal{C}^\times\cap\mathcal{C}_2$, then it belongs to $\mathcal{C}^\times\cap\mathcal{C}_1$, hence to $\mathcal{C}_1^\times$ since $\mathcal{C}_1$ is a head-subcategory of $\mathcal{C}$. So ϵ belongs to $\mathcal{C}_1^\times\cap\mathcal{C}_2$, hence to $\mathcal{C}_2 i$ since $\mathcal{C}_2$ is a head-subcategory of $\mathcal{C}_1$. So $\mathcal{C}_2$ is closed under inverse in $\mathcal{C}$. Next assume $g\in\mathcal{C}$. Let g' be a $\mathcal{C}_1$-head of g in $\mathcal{C}$, and g'' be a $\mathcal{C}_2$-head of g' in $\mathcal{C}_1$. Assume $h\in\mathcal{C}_2$ and $h\preccurlyeq g$. As h belongs to $\mathcal{C}_1$, we must have $h\preccurlyeq g'$. Then, as h belongs to $\mathcal{C}_2$, we must have $h\preccurlyeq g''$. So g'' is a $\mathcal{C}_2$-head of g, and $\mathcal{C}_2$ is a head-subcategory of $\mathcal{C}$.

Exercice 82 (**recognizing compatible IV**).— *Assume that $\mathcal{S}$ is a Garside family in a left-cancellative category $\mathcal{C}$ and $\mathcal{C}_1$ is a subcategory of $\mathcal{C}$ that is closed under right-quotient in $\mathcal{C}$. Show that $\mathcal{C}_1$ is compatible with $\mathcal{S}$ if and only if, putting $\mathcal{S}_1 = \mathcal{S}\cap\mathcal{C}_1$,* (i) *the family $\mathcal{S}_1$ is a Garside family in $\mathcal{C}_1$, and* (ii) *a $\mathcal{C}_1$-path is strictly $\mathcal{S}_1$-normal in $\mathcal{C}_1$ if and only if it is strictly $\mathcal{S}$-normal in $\mathcal{C}$.*

Solution. Assume that $\mathcal{C}_1$ is compatible with $\mathcal{S}$. By definition, $\mathcal{S}_1$ is a Garside family in $\mathcal{C}_1$, so (i) is satisfied. Next, assume that $g_1|\cdots|g_q$ is a strict $\mathcal{S}$-normal $\mathcal{C}_1$-path. By assumption, $g_1|\cdots|g_q$ is an $\mathcal{S}_1$-normal path in $\mathcal{C}_1$. Moreover, by definition, $g_1, ..., g_q$ are not invertible in $\mathcal{C}$, hence they are not invertible either in $\mathcal{C}_1$, and $g_1, ..., g_{q-1}$ belong to $\mathcal{S}$ and to $\mathcal{C}_1$, hence to $\mathcal{S}_1$. So $g_1|\cdots|g_q$ is strictly $\mathcal{S}_1$-normal in $\mathcal{C}_1$. Conversely, assume that $g_1|\cdots|g_q$ is strictly $\mathcal{S}_1$-normal in $\mathcal{C}_1$. As $\mathcal{C}_1$ is compatible with $\mathcal{S}$, the path $g_1|\cdots|g_q$ is $\mathcal{S}$-normal in $\mathcal{C}$. Moreover, the assumption that $g_1, ..., g_{q-1}$ belong to $\mathcal{S}_1$ implies that they belong to $\mathcal{S}$. Finally, the assumption that no g_i is invertible in $\mathcal{C}_1$ implies that they are not invertible in $\mathcal{C}$ either. So (ii) is satisfied.

Conversely, assume that $\mathcal{C}_1$ satisfies (i) and (ii). Assume that $g_1|g_2$ is an $\mathcal{S}_1$-path. Then g_1g_2 is invertible in $\mathcal{C}_1$ if and only if it is invertible in $\mathcal{C}$. In this case, $g_1|g_2$ is both $\mathcal{S}_1$-normal in $\mathcal{C}_1$ and $\mathcal{S}$-normal in $\mathcal{C}$. Otherwise, if g_1g_2 belongs to $\mathcal{S}_1\mathcal{C}_1^\times$, then $g_1|g_2$ is $\mathcal{S}_1$-normal in $\mathcal{C}_1$ if and only if g_2 is invertible in $\mathcal{C}_1$ and, similarly, it is $\mathcal{S}$-normal in $\mathcal{C}$ if and only if g_2 is invertible in $\mathcal{C}$, so both conditions are equivalent. Finally, if g_1g_2 has $\mathcal{S}_1$-length 2, it admits a strict $\mathcal{S}_1$-normal decomposition in $\mathcal{C}_1$, say $h_1|h_2$. Then $g_1|g_2$ is $\mathcal{S}_1$-normal in $\mathcal{C}_1$ if and only if there exist ϵ invertible in $\mathcal{C}_1$ satisfying $g_1 = h_1\epsilon$ and $h_2 = \epsilon g_2$; similarly, $g_1|g_2$ is $\mathcal{S}$-normal in $\mathcal{C}$ if and only if there exist ϵ invertible in $\mathcal{C}$ satisfying $g_1 = h_1\epsilon$ and $h_2 = \epsilon g_2$: both conditions are equivalent again. So an $\mathcal{S}_1$-path is $\mathcal{S}_1$-normal in $\mathcal{C}_1$ if and only if it is $\mathcal{S}$-normal in $\mathcal{C}$, and, by definition, $\mathcal{C}_1$ is compatible with $\mathcal{S}$.

Exercice 83 (**inverse image**).— *Assume that $\mathcal{C}, \mathcal{C}'$ are left-cancellative categories, ϕ is a functor from $\mathcal{C}$ to $\mathcal{C}'$, and $\mathcal{C}'_1$ is a subcategory of $\mathcal{C}'$ that is closed under left- and right-divisor. Show that the subcategory $\phi^{-1}(\mathcal{C}'_1)$ is compatible with every Garside family of $\mathcal{C}$.* (ii) *Let B^+ be the Artin–Tits monoid of type B as defined in Example VII.4.21. Show that the map ϕ defined by $\phi(\sigma_0) = 1$ and $\phi(\sigma_i) = 0$ for $i \geqslant 1$ extends into a homomorphism of B^+ to $\mathbb{N}$, and that the submonoid $N = \{g \in M \mid \phi(g) = 0\}$ of B^+ is compatible with every Garside family of B^+.*

Solution. (i) By Proposition VII.1.18, the subcategory $\phi^{-1}(\mathcal{C}'_1)$ is closed under right-quotient and under $=^\times$. Then apply Proposition VII.2.21. (ii) Use (i)

Exercice 84 (**intersection**).— *Assume that $\mathcal{S}$ is a Garside family in a left-cancellative category $\mathcal{C}$.* (i) *Let $\mathcal{F}$ be the family of all subcategories of $\mathcal{C}$ that are closed under right-quotient, compatible with $\mathcal{S}$, and $=^\times$-closed. Show that every intersection of elements of $\mathcal{F}$ belongs to $\mathcal{F}$.* (ii) *Same question when "$=^\times$-closed" is replaced with "including $\mathcal{C}^\times$".* (iii) *Same question when $\mathcal{C}$ contains no nontrivial invertible element and "$=^\times$-closed" is skipped.*

Solution. (i) Let $(\mathcal{C}_i)_{i\in I}$ be a family of $=^\times$-closed subcategories of $\mathcal{C}$ that are compatible with $\mathcal{S}$. First, an intersection of subcategories is a subcategory. Next, an intersection of $=^\times$-closed families is $=^\times$-closed and, similarly, an intersection of families that are closed under right-quotient is closed under right-quotient. Finally, let g belong to $\bigcap \mathcal{C}_i$. Then g admits an $\mathcal{S}$-normal decomposition $s_1|\cdots|s_p$. By Lemma VII.2.20, the latter is $(\mathcal{S} \cap \mathcal{C}_i)$-normal for every i, hence it its entries lie in every subfamily $\mathcal{C}_i$, hence in their intersection. By Proposition VII.2.21 we deduce that $\bigcap \mathcal{C}_i$ is compatible with $\mathcal{S}$.

(ii) An intersection of families that include $\mathcal{C}^\times$ includes $\mathcal{C}^\times$. On the other hand, a subcategory that includes $\mathcal{C}^\times$ is $=^\times$-closed and we apply (i).

(iii) In this case, every subcategory is $=^{\times}$-closed, and we apply (i).

Exercice 85 (**fixed points**).— *Assume that $\mathcal{C}$ is a left-cancellative category and $\phi : \mathcal{C} \to \mathcal{C}$ is a functor. Show that the fixed point subcategory $\mathcal{C}^\phi$ is compatible with $\mathcal{C}$ viewed as a Garside family in itself.*

Solution. First, assume that ϵ belongs to $\mathcal{C}(x,y) \cap \mathcal{C}^\phi$. Necessarily we have $\phi(x) = x$. As ϕ is a functor, we find $\epsilon\epsilon^{-1} = 1_x = \phi(1_x) = \phi(\epsilon\epsilon^{-1}) = \phi(\epsilon)\phi(\epsilon^{-1}) = \epsilon\phi(\epsilon^{-1})$, whence $\epsilon^{-1} = \phi(\epsilon^{-1})$, so $\mathcal{C}^\times \cap \mathcal{C}^\phi \subseteq (\mathcal{C}^\phi)^\times$ holds. Next, we have $\mathcal{C}^\sharp = \mathcal{C}$, so $\mathcal{C}^\sharp \cap \mathcal{C}^\phi \subseteq \mathcal{S}^\phi$ trivially holds, and $\mathcal{C}^\phi$ and $\mathcal{C}$ satisfy (the counterpart of) (VII.2.11). On the other hand, every element g of $\mathcal{C}$ admits the $\mathcal{C}$-normal decomposition g, so, in particular, every element of $\mathcal{C}^\phi$ has a $\mathcal{C}$-normal decomposition whose entries lie in $\mathcal{C}^\phi$. So (the counterpart of) (VII.2.12) is satisfied and, by Proposition VII.2.10, $\mathcal{C}^\phi$ is compatible with $\mathcal{C}$.

Exercice 86 (**connection between closure properties**).— *Assume that $\mathcal{S}$ is a subfamily in a left-cancellative category $\mathcal{C}$ and $\mathcal{S}_1$ is a subfamily of $\mathcal{S}$.* (i) *Show that, if $\mathcal{S}_1$ is closed under product, inverse, and right-complement in $\mathcal{S}$, then $\mathcal{S}_1$ is closed under right-quotient in $\mathcal{S}$.* (ii) *Assume that $\mathcal{S}_1$ is closed under product and right-complement in $\mathcal{S}$. Show that, if $\mathcal{S}$ is closed under left-divisor in $\mathcal{C}$, then $\mathcal{S}_1$ is closed under right-comultiple in $\mathcal{S}$. Show that, if $\mathcal{S}$ is closed under right-diamond in $\mathcal{C}$, then $\mathcal{S}ub(\mathcal{S}_1)$ is closed under right-comultiple in $\mathcal{S}$.* (iii) *Show that, if $\mathcal{S}_1$ is closed under identity and product in $\mathcal{S}$, then $\mathcal{S}_1$ is closed under inverse and right-diamond in $\mathcal{S}$ if and only if $\mathcal{S}_1$ is closed under right-quotient and right-comultiple in $\mathcal{S}$.*

Solution. (i) The argument is the same as for Lemma VII.1.8. Assume $sg = t$ with s, t in $\mathcal{S}_1$ and g in $\mathcal{S}$. As $\mathcal{S}_1$ is closed under right-complement in $\mathcal{S}$, there exist s', t' in $\mathcal{S}_1$ and r in $\mathcal{S}$ satisfying $st' = ts'$, $1_y = s'r$, and $g = t'r$, where y is the target of t. The second equality implies that s' is invertible, and the assumption that $\mathcal{S}_1$ is closed under inverse in $\mathcal{S}$ then implies that s'^{-1}, that is, r, lies in $\mathcal{S}_1$. Hence $t'r$, that is g, belongs to $\mathcal{S}_1^{[2]} \cap \mathcal{S}$, hence to $\mathcal{S}_1$ as the latter is closed under product in $\mathcal{S}$. So $\mathcal{S}_1$ is closed under right-quotient in $\mathcal{S}$.

(iii) Assume that $\mathcal{S}_1$ is closed under inverse and right-diamond in $\mathcal{S}$. Then, by definition, $\mathcal{S}_1$ is closed under right-comultiple and right-complement, and, by Exercice 89 (transfer of closure), it is closed under right-quotient.

Conversely, assume that $\mathcal{S}_1$ is closed under right-quotient and right-comultiple in $\mathcal{S}$. First, the assumption that $\mathcal{S}_1$ is closed under right-quotient trivially implies that $\mathcal{S}_1$ is closed under inverse. Next, by the same argument as in Lemma IV.1.8, the assumption that $\mathcal{S}_1$ is closed under right-comultiple and right-quotient implies that it is closed under right-diamond.

Exercice 87 (**subgerm**).— *Assume that $\underline{\mathcal{S}}$ is a left-cancellative germ and $\underline{\mathcal{S}}_1$ is a subgerm of $\mathcal{S}$ such that the relation $\preccurlyeq_{\mathcal{S}_1}$ is the restriction to $\mathcal{S}_1$ of the relation $\preccurlyeq_{\mathcal{S}}$. Show that $\mathcal{S}_1$ is closed under right-quotient in $\mathcal{S}$.*

Solution. Assume $f \bullet h = g$ with $f, g \in \mathcal{S}_1$. By definition, $f \preccurlyeq_{\mathcal{S}} g$ holds, hence so does $f \preccurlyeq_{\mathcal{S}_1} g$. This means that there exists h_1 in $\mathcal{S}_1$ satisfying $f \bullet h_1 = g$. The assumption that $\mathcal{S}$ is left-cancellative implies $h_1 = h$, whence $h \in \mathcal{S}$. So $\mathcal{S}_1$ is closed under right-quotient in $\mathcal{S}$.

Exercice 88 (**transitivity of closure**).— *Assume that $\underline{\mathcal{S}}_1$ is a subgerm of a Garside germ $\underline{\mathcal{S}}$, the subcategory $\mathcal{S}ub(\mathcal{S}_1)$ is closed under right-quotient and right-diamond in $\mathcal{C}at(\underline{\mathcal{S}})$, and $\mathcal{S}_1$ is closed under inverse and right-complement (resp. right-diamond) in $\mathcal{S}ub(\mathcal{S}_1)$. Show that $\mathcal{S}_1$ is closed under inverse and right-complement (resp. right-diamond) in $\mathcal{S}$.*

Solution. By Lemma VII.1.7, the assumption that $\mathcal{S}ub(\mathcal{S}_1)$ is closed under right-quotient implies that it is closed under inverse, and Lemma VII.3.7, which is valid since the assumption that $\underline{\mathcal{S}}$ is a Garside germ implies that $\mathcal{S}$ is closed under right-quotient in $\mathcal{C}at(\underline{\mathcal{S}})$, then implies that $\mathcal{S}_1$ is closed under inverse in $\mathcal{S}$. Next, by Lemma VII.3.8 applied with $\mathcal{S}ub(\mathcal{S}_1)$ in place of $\mathcal{S}$, the assumption that $\mathcal{S}_1$ is closed under right-complement (*resp.* right-diamond) in $\mathcal{S}ub(\mathcal{S}_1)$ implies that $\mathcal{S}_1$ is closed under right-complement (*resp.* right-diamond) in $\mathcal{C}at(\underline{\mathcal{S}})$. Now, as $\mathcal{S}$ is a solid Garside family in $\mathcal{C}at(\underline{\mathcal{S}})$, it is closed under right-divisor in $\mathcal{C}at(\underline{\mathcal{S}})$, hence a fortiori under right-quotient. Applying once more Lemma VII.3.8, we deduce from the fact that $\mathcal{S}_1$ is closed under right-complement (*resp.* right-diamond) in $\mathcal{C}at(\underline{\mathcal{S}})$ that $\mathcal{S}_1$ is closed under right-complement (*resp.* right-diamond) in $\mathcal{S}$.

Exercice 89 (**transfer of closure**).— *Assume that $\mathcal{C}$ is a left-cancellative category, $\mathcal{S}$ is a subfamily of $\mathcal{C}$, and $\mathcal{S}_1$ is a subfamily of $\mathcal{S}$ that is closed under identity and product.* (i) *Show that, if $\mathcal{S}ub(\mathcal{S}_1)$ is closed under right-quotient in $\mathcal{C}$, then $\mathcal{S}_1$ is closed under right-quotient in $\mathcal{S}$.* (ii) *Show that, if, moreover, $\mathcal{S}$ is closed under right-quotient in $\mathcal{C}$, then $\mathcal{S}_1$ is closed under right-quotient in $\mathcal{S}ub(\mathcal{S}_1)$.*

Solution. (i) Assume that t belongs to $\mathcal{S}$ and s and st belong to $\mathcal{S}_1$. Then s and st belong to $\mathcal{S}ub(\mathcal{S}_1)$, so, as the latter is assumed to be closed under right-quotient, t must belong to $\mathcal{S}ub(\mathcal{S}_1)$, hence to $\mathcal{S}ub(\mathcal{S}_1) \cap \mathcal{S}$. By Proposition VII.3.3, the latter is $\mathcal{S}_1$. So $\mathcal{S}_1$ is closed under right-quotient in $\mathcal{S}$.

(ii) Assume that $\mathcal{S}$ is closed under right-quotient in $\mathcal{C}$. Assume that t belongs to $\mathcal{S}ub(\mathcal{S}_1)$ and s and st belong to $\mathcal{S}_1$. Then s and st belong to $\mathcal{S}$, so the assumption implies that t belong to $\mathcal{S}$, hence to $\mathcal{S}ub(\mathcal{S}_1) \cap \mathcal{S}$, which is $\mathcal{S}_1$ by Proposition VII.3.3. So $\mathcal{S}_1$ is closed under right-quotient in $\mathcal{S}ub(\mathcal{S}_1)$.

Exercice 90 (**braid subgerm**).— *Let $\underline{\mathcal{S}}$ be the six-element Garside germ associated with the divisors of Δ_3 in the braid monoid B_3^+.* (i) *Describe the subgerm $\underline{\mathcal{S}}_1$ of $\underline{\mathcal{S}}$ generated by σ_1 and σ_2. Compare $\mathcal{M}on(\underline{\mathcal{S}}_1)$ and $\mathcal{S}ub(\mathcal{S}_1)$ (describe them explicitly).* (ii) *Same questions with σ_1 and $\sigma_2\sigma_1$. Is $\mathcal{S}_1$ closed under right-quotient in $\underline{\mathcal{S}}$ in this case?*

Solution. (i) The subgerm of $\underline{\mathcal{S}}$ generated by σ_1 and σ_2 is the closure of $\{\sigma_1, \sigma_2\}$ under identity and product in $\underline{\mathcal{S}}$: so it contains 1, and, as $\sigma_1 \bullet \sigma_2$ is defined in $\underline{\mathcal{S}}$, it contains $\sigma_1 \bullet \sigma_2$, that is, $\sigma_1\sigma_2$. Then it contains $\sigma_1\sigma_2 \bullet \sigma_1$, which is Δ_3, etc. Finally, one finds $\underline{\mathcal{S}}_1 = \underline{\mathcal{S}}$, whence $\mathcal{M}on(\underline{\mathcal{S}}_1) = \mathcal{S}ub(\mathcal{S}_1) = B_3^+$.

(ii) The closure of $\{\sigma_1, \sigma_2\sigma_1\}$ under identity and product is $\{1, \sigma_1, \sigma_2\sigma_1, \Delta_3\}$. In the associated germ $\underline{\mathcal{S}}_1$, the only nontrivial product is $\sigma_1 \bullet_1 \sigma_2\sigma_1 = \Delta_3$, so $\mathcal{M}on(\underline{\mathcal{S}}_1)$ is $\langle \mathtt{a}, \mathtt{b}, \Delta_3 \,|\, \mathtt{ab} = \Delta_3 \rangle^+$, that is, a free monoid based on $\mathtt{a}$ and $\mathtt{b}$. On the other hand, in $\mathcal{S}ub(\mathcal{S}_1)$, the relation $(\sigma_2\sigma_1)^3 = \Delta_3^2$ holds, corresponding to $\mathtt{b}^3 = (\mathtt{ab})^2$, which fails in $\mathcal{M}on(\underline{\mathcal{S}}_1)$: so $\mathcal{M}on(\underline{\mathcal{S}}_1)$ is not isomorphic to $\mathcal{S}ub(\mathcal{S}_1)$. Here $\mathcal{S}ub(\mathcal{S}_1)$ is not closed under right-quotient in $\underline{\mathcal{S}}$, as Δ_3 and $\sigma_2\sigma_1$ lie in $\mathcal{S}ub(\mathcal{S}_1)$, but we have $\Delta_3 = \sigma_2\sigma_1 \bullet \sigma_2$ in $\underline{\mathcal{S}}$ and $\sigma_2 \notin \mathcal{S}_1$.

Exercice 91 ($=^\times$**-closed**).— *Show that, if $\underline{\mathcal{S}}_1$ is a subgerm of an associative germ $\underline{\mathcal{S}}$, then $\mathcal{S}ub(\mathcal{S}_1)$ is $=^\times$-closed in $\mathcal{C}at(\underline{\mathcal{S}})$ if and only if $\underline{\mathcal{S}}_1$ is $=^\times_{\mathcal{S}}$-closed in $\mathcal{S}$.*

Solution. Assume that $\underline{\mathcal{S}}_1$ is an $=^\times_{\mathcal{S}}$-closed subgerm of $\underline{\mathcal{S}}$. Assume that g is an element of $\mathcal{S}ub(\mathcal{S}_1)$ and $g' =^\times g$ holds in $\mathcal{C}at(\underline{\mathcal{S}})$. By definition, g admits an $\mathcal{S}_1$-decomposition, say $s_1|\cdots|s_p$, and, in $\mathcal{C}at(\underline{\mathcal{S}})$, we then have $g' = s_1\cdots s_p\epsilon$. As $\mathcal{S}$ is closed under left-divisor in $\mathcal{C}at(\underline{\mathcal{S}})$ since $\underline{\mathcal{S}}$ is right-associative, the assumption that s_p lies in $\mathcal{S}$ implies that its left-divisor $s_p\epsilon$ also lies in $\mathcal{S}$, that is, $s_p \bullet \epsilon$ is defined in $\underline{\mathcal{S}}$. The assumption that $\underline{\mathcal{S}}_1$ is $=^\times_{\mathcal{S}}$-closed in $\underline{\mathcal{S}}$ then implies that $s_p \bullet \epsilon$ belongs to $\mathcal{S}_1$, which implies that g', which is $s_1\cdots s_{p-1}(s_p\epsilon)$, lies in $\mathcal{S}ub(\mathcal{S}_1)$. Conversely, assume that $\mathcal{S}ub(\mathcal{S}_1)$ is an $=^\times$-closed subcategory of $\mathcal{C}at(\underline{\mathcal{S}})$. Assume that s belongs to $\mathcal{S}_1$ and $s' =^\times_{\mathcal{S}} s$ holds in $\underline{\mathcal{S}}$. This means that there exists ϵ in $\mathcal{S}^\times$ satisfying $s' = s \bullet \epsilon$. Then, in $\mathcal{C}at(\underline{\mathcal{S}})$, we have $s' = s\epsilon$, whence $s' =^\times s$. The assumption that $\mathcal{S}ub(\mathcal{S}_1)$ is $=^\times$-closed implies $s' \in \mathcal{S}ub(\mathcal{S}_1)$, whence $s' \in \mathcal{S}ub(\mathcal{S}_1) \cap \mathcal{S}$. As $\underline{\mathcal{S}}_1$ is a subgerm of $\underline{\mathcal{S}}$, the latter family is $\mathcal{S}_1$.

Exercice 92 (**correct** *vs.* **mcms**).— *Assume that $\mathcal{C}$ and $\mathcal{C}'$ are left-cancellative categories and $\mathcal{S}$ is included in $\mathcal{C}$. Assume moreover that $\mathcal{C}$ and $\mathcal{C}'$ admit right-mcms and $\mathcal{S}$ is closed under right-mcm. Show that a functor ϕ from $\mathcal{C}$ to $\mathcal{C}'$ is correct for right-comultiples on $\mathcal{S}$ if and only if, for all s, t in $\mathcal{S}$, every right-mcm of $\phi(s)$ and $\phi(t)$ is $=^\times$-equivalent to the image under ϕ of a right-mcm of s and t.*

Solution. Assume that ϕ is correct for right-comultiples on $\mathcal{S}$, that s, t lie in $\mathcal{S}$, and that h is a right-mcm of $\phi(s)$ and $\phi(t)$. By definition, there exists a common right-multiple r of s and t that lies in $\mathcal{S}$ and satisfies $\phi(r) \preccurlyeq h$. As $\mathcal{C}$ admits right-mcms, there exists a right-mcm r' of s and t satisfying $r' \preccurlyeq r$ and, as $\mathcal{S}$ is closed under right-mcm, r' lies in $\mathcal{S}$. Then $\phi(r') \preccurlyeq h$ holds, and $\phi(r')$ is a common right-multiple of $\phi(s)$ and $\phi(t)$. As r is minimal, we deduce $\phi(r') =^\times h$. So the condition is necessary. Conversely, assume that ϕ preserves mcms in the sense of the statement. Assume that s, t lie in $\mathcal{S}$, and h is a common right-multiple of $\phi(s)$ and $\phi(t)$. As $\mathcal{C}'$ admits right-mcms, h is a right-multiple of some right-mcm of $\phi(s)$ and $\phi(t)$, hence, by (ii), of some element $\phi(r)$ where r is a right-mcm of s and t. By assumption, r belongs to $\mathcal{S}$, and there exist s', t' in $\mathcal{C}$ satisfying $st' = ts' = r$. Hence ϕ is correct for right-comultiples on $\mathcal{S}$.

Chapter VIII: Conjugacy

Skipped proofs

(none)

Solution to selected exercises

Exercice 97 (**quasi-distance**).— (i) *In Context VIII.3.7, show that $\ell_\Delta(g) = \sup_\Delta(g^0)$ holds for every g in $\mathcal{G}$.* (ii) *For g, g' in $\mathcal{G}$, define* $\mathrm{dist}(g, g')$ *to be ∞ if g, g' do not share the same source, and to be $\ell_\Delta(g^{-1}g')$ otherwise. Show that* dist *is a quasi-distance on $\mathcal{G}$ that is compatible with $=_\Delta$.*

Solution. (ii) The canonical length is invariant under left- and right-multiplication by Δ and, therefore, dist takes constant values on $=_\Delta$-classes.

Chapter IX: Braids

Skipped proofs

(none)

Solution to selected exercises

Exercice 102 (**smallest Garside, right-angled type**).— *Assume that B^+ is a right-angled Artin–Tits monoid, that is, B^+ is associated with a Coxeter system (W,Σ) satisfying $m_{s,t}\in\{2,\infty\}$ for all s,t in Σ.* (i) *For $I\subseteq\Sigma$, denote by Δ_I the right-lcm (here the product) of the elements of I, when it exists (that is, when the elements pairwise commute). Show that the divisors of Δ_I are the elements Δ_J with $J\subseteq I$.* (ii) *Deduce that the smallest Garside family in B^+ is finite and consists of the elements Δ_I for $I\subseteq\Sigma$.*

Solution. (i) Clearly $J\subseteq I$ implies $\Delta_J\preccurlyeq\Delta_I$ and $\Delta_J\widetilde{\preccurlyeq}\Delta_I$, since the elements of I commute. Conversely, the point is to see that, if u is a word in S that involves at least one letter not in I, then (the class of) u cannot right-divide Δ_I: this is so since, when left-reversing $\Delta_I\overline{u}$, a negative letter $\overline{s}$ can only vanish when it is adjacent to the positive letter s, and it, if u is a word in I in which some letter is repeated twice, then the second occurrence cannot vanish as we are arguing in the free abelian monoid based on I.

(ii) Let S be the family of all elements Δ_I with I a set of pairwise commuting atoms in B^+. As the smallest Garside family of B^+ includes I and is closed under right-lcm, it must include S. On the other hand, S includes Σ, it is closed under right-lcm by definition, and it is closed under right-divisor by (i). Corollary IV.2.29 (recognizing Garside, right-lcm case) implies that S is a Garside family in B^+.

Exercice 103 (**smallest Garside, large type**).— *Assume that B^+ is an Artin–Tits monoid of large type, B^+ is associated with a Coxeter system (W,Σ) satisfying $m_{s,t}\geqslant 3$ for all s,t in Σ. Put*

$\Sigma_1=\{s\in\Sigma\mid\forall r\in\Sigma\,(m_{r,s}=\infty)\}$,

$\Sigma_2=\{(s,t)\in\Sigma^2\mid m_{s,t}<\infty$ *and* $\forall r\in\Sigma\,(m_{r,s}+m_{r,t}=\infty\}$,

$\Sigma_3=\{(r,s,t)\in\Sigma^3\mid m_{r,s}+m_{r,t}+m_{s,t}<\infty\}$,

and $E=\Sigma_1\cup\{\Delta_{s,t}\mid(s,t)\in\Sigma_2\}\cup\{r\Delta_{s,t}\mid(r,s,t)\in\Sigma_3\}$ (we write $\Delta_{s,t}$ for the right-lcm of s and t when it exists). (i) *Explicitly describe the elements of the closure S of E under right-divisor, and deduce that S is finite.* (ii) *Show that S is closed under right-lcm and deduce that S is a Garside family in B^+.* (iii) *Show that S is the smallest Garside family in B^+.* (iv) *Show that, if Σ has n elements and $m_{s,t}\neq\infty$ holds for all s,t, then E has $3\binom{n}{3}$ elements.* (v) *Show that, if Σ has n elements and $m_{s,t}=m$ holds for all s,t, then S has $(n+2m-5)\binom{n}{2}+n+1$ elements. Apply to $n=m=3$.*

Solution. (i) For s in Σ_1, the only right-divisors of s are 1 and s; for (s,t) in Σ_2, the right-divisors of $\Delta_{s,t}$ are the elements $(s|t)^{[k]}$ and $(t|s)^{[k]}$ with $k \leqslant m_{s,t}$: this is so because right-divisors are detected using left-reversing (Section II.4) and it is clear that, if u is a word involving a letter different from s and t or if u involves two letters s or two letters t, or if u is of the form $(s|t)^{[k]}$ and $(t|s)^{[k]}$ with $k > m_{s,t}$, then left-reversing $\Delta_{s,t}u^{-1}$ cannot result in a positive word. Finally, for (r,s,t) in Σ_3, the right-divisors of $r\Delta_{s,t}$ are $r\Delta_{s,t}$ and the elements $(s|t)^{[k]}$ and $(t|s)^{[k]}$ with $k \leqslant m_{s,t}$: again, this is so because left-reversing $\Delta_{s,t}u^{-1}$ can result in a positive word only in the cases described above, and it can result in r^{-1} concatenated with a positive word only if u is $r\Delta_{s,t}$ itself.

(ii) We have to analyze when two elements of S may admit a common right-multiple. As there are three cases for each factor, nine cases are to be considered. Many of them are trivial. First, all cases involving an atom of Σ_1 are trivial. Then, in the case of two elements $(s|t)^{[k]}$ and $(s'|t')^{[k']}$, the case $\{s,t\} = \{s',t'\}$ is obvious. Otherwise, for $s' = s$ and $t' \neq t$, the only cases when a common right-multiple may exist are the trivial cases $k = 1$ or $k' = 1$, plus the case $k = k' = 2$, in which case there exists a common right-multiple for $m_{t,t'} < \infty$, and the right-lcm is then $s\Delta_{t,t'}$, an element of S. Finally, for $\#\{s,t,s',t'\} = 4$, no common right-multiple may exist for $k, k' \geqslant 2$, and the remaining cases are treated as above. The cases of an element $r\Delta_{s,t}$ and an element $(s'|t')^{[k']}$ and of two elements $r\Delta_{s,t}$ and $r'\Delta_{s',t'}$ are treated in the same way: once again, the point is that common right-multiples may exist only in the obvious cases. Finally, S is closed under right-lcm. As, by definition, S includes Σ and is closed under right-divisor, Corollary IV.2.29 (recognizing Garside, right-lcm case) implies that S is a Garside family in B^+.

(iii) Conversely, every Garside family S' of B^+ containing 1 must include S. Indeed, S' must include Σ, hence all elements $\Delta_{s,t}$ since it is closed under right-lcm. Moreover, for (r,s,t) in Σ_3, the family S' must contain rs since it is closed under right-divisor and rs right-divides $\Delta_{r,s}$; it contains rt for a similar reason and, therefore, it contains the right-lcm $r\Delta_{s,t}$ of rs and rt. So S' includes E, hence S, and S is the smallest Garside family containing 1 in B^+.

(iv) When no coefficient $m_{s,t}$ is ∞, Σ_1 and Σ_2 are empty and E consists of all elements $r\Delta_{s,t}$ with r,s,t pairwise distinct atoms in B^+. As $m_{s,t}$ does not matter, there exist $3\binom{n}{3}$ such elements for Σ of size n.

(v) By the description of (i), S comprises 1, plus the n atoms, plus, for each of the $\binom{n}{2}$ pairs (s,t), the $2m-3$ right-divisors of $\Delta_{s,t}$ of length $\geqslant 2$, plus, for each of the $\binom{n}{3}$ triples (r,s,t), the 3 elements $r\Delta_{s,t}, s\Delta_{t,r}, t\Delta_{r,s}$, whence the formula. The case $n = m = 3$ corresponds to the affine type $\tilde{\mathrm{A}}_2$, and confirms that the smallest Garside family containing 1 has 16 elements, as seen in Reference Structure 9, page 111.

Chapter X: Deligne–Lusztig varieties

Skipped proofs

(none)

Solution to selected exercises

(none)

Chapter XI: Left self-distributivity

Skipped proofs

Lemma XI.1.8.— *Assume that F is a partial action of a monoid M on a set X.*
(i) *If the monoid M is left-cancellative, then so is the category $\mathcal{C}(M,F)$.*
(ii) *Conversely, if F is proper and $\mathcal{C}(M,F)$ is left-cancellative, then so is M.*

Proof. (i) Assume that M is left-cancellative and, in $\mathcal{C}(M,F)$, the equality $x \xrightarrow{g} y \cdot y \xrightarrow{h} z = x \xrightarrow{g} y \cdot y \xrightarrow{h'} z'$ holds. This implies $x \xrightarrow{gh} z = x \xrightarrow{gh'} z'$, whence, by definition, $gh = gh'$ and $z = z'$, and $h = h'$ since M is left-cancellative. We deduce $y \xrightarrow{h} z = y \xrightarrow{h'} z'$, and $\mathcal{C}(M,F)$ is left-cancellative.

(ii) Conversely, assume that F is proper and $\mathcal{C}(M,F)$ is left-cancellative. Assume $gh = gh'$. As F is proper, there exists x in X such that both $x \bullet gh$ and $x \bullet gh'$ are defined. Let $z = x \bullet gh$. By (XI.1.3), $x \bullet g$ must be defined and, putting $y = x \bullet g$, we have $z = y \bullet h = y \bullet h'$. Then, in $\mathcal{C}(M,F)$, we have $x \xrightarrow{g} y \cdot y \xrightarrow{h} z = x \xrightarrow{g} y \cdot y \xrightarrow{h'} z'$. As $\mathcal{C}(M,F)$ is left-cancellative, we deduce $y \xrightarrow{h} z = y \xrightarrow{h'} z'$, whence $h = h'$. Hence M is left-cancellative. □

Lemma XI.1.9.— *Assume that F is a partial action of a monoid M on a set X.*

(i) *Assume that $x \bullet g$ is defined. Then $y \xrightarrow{h} \text{-} \preccurlyeq x \xrightarrow{g} \text{-}$ holds in $\mathcal{C}(M,F)$ if and only if we have $y = x$ and $h \preccurlyeq g$ in M.*

(ii) *Assume that $x \bullet f$ and $x \bullet g$ are defined. Then $x \xrightarrow{h} \text{-}$ is a left-gcd of $x \xrightarrow{f} \text{-}$ and $x \xrightarrow{g} \text{-}$ in $\mathcal{C}(M,F)$ if and only if h is a left-gcd of f and g in M.*

Proof. (i) Assume $y \xrightarrow{h} y \bullet h \cdot x' \xrightarrow{g'} \text{-} = (x, g, \text{-})$. Then we have $y = x$ and $hg' = g$, hence $h \preccurlyeq g$. Conversely, assume $h \preccurlyeq g$, say $hg' = g$. By (XI.1.3), the assumption that $x \bullet g$ is defined implies that $x \bullet h$ is defined, say $x \bullet h = x'$. Then we have $x \xrightarrow{h} x' \cdot x' \xrightarrow{g'} \text{-} = x \xrightarrow{g} \text{-}$, whence $x \xrightarrow{h} \text{-} \preccurlyeq x \xrightarrow{g} \text{-}$ in $\mathcal{C}(M,F)$.

(ii) Assume that $x \xrightarrow{h} \text{-}$ is a left-gcd of $x \xrightarrow{f} \text{-}$ and $x \xrightarrow{g} \text{-}$ in $\mathcal{C}(M,F)$. By (i), h left-divides f and g in M. Let h' be a common left-divisor of f and g in M. By (XI.1.3), the assumption that $x \bullet f$ is defined implies that $x \bullet h'$ is defined. Then $x \xrightarrow{h} \text{-}$ is a common left-divisor of $x \xrightarrow{f} \text{-}$ and $x \xrightarrow{g} \text{-}$ in $\mathcal{C}(M,F)$, hence it is a left-divisor of $x \xrightarrow{h} \text{-}$. By (i), this implies $h' \preccurlyeq h$. So h is a left-gcd of f and g.

Conversely, assume that h is a left-gcd of f and g. By (i), $x \xrightarrow{h} \text{-}$ left-divides $x \xrightarrow{f} \text{-}$ and $x \xrightarrow{g} \text{-}$ in $\mathcal{C}(M,F)$. Now, consider an arbitrary common left-divisor of $\mathtt{NF}(x, f, \text{-})$ and $x \xrightarrow{g} \text{-}$. By (i), it must be of the form $x \xrightarrow{h'} \text{-}$ with h' left-dividing f and g. Then h left-divides h in M, and $x \xrightarrow{h'} \text{-}$ left-divides $x \xrightarrow{h} \text{-}$ in $\mathcal{C}(M,F)$. □

Lemma XI.4.16.— *Assume that M is a left-cancellative monoid, F is a proper partial action of M on some set X, and $(\Delta_x)_{x\in X}$ is a right-Garside map on M with respect to F. Assume moreover that S is a family of atoms that generate M. Now assume that $\pi : M \to \underline{M}$ is a surjective homomorphism, $\pi_\bullet : X \to \underline{X}$ is a surjective map, and, for all x in X and g in M,*

(XI.4.17) *The value of $\pi_\bullet(x \bullet g)$ only depends on $\pi_\bullet(x)$ and $\pi(g)$;*

(XI.4.18) *The value of $\pi(\Delta_x)$ only depends on $\pi_\bullet(x)$.*

Assume finally that $\widetilde{\pi} : \underline{S} \to S$ is a section of π such that, for x in X, $\underline{s}$ in $\underline{S}$, and $\underline{w}$ in $\underline{S}^*$,

(XI.4.19) *If $\pi_\bullet(x) \bullet \underline{s}$ is defined, then so is $x \bullet \widetilde{\pi}(\underline{s})$.*

(XI.4.20) *The relation $[\underline{w}] \preccurlyeq \Delta_{\pi_\bullet(x)}$ implies $[\widetilde{\pi}^*(\underline{w})] \preccurlyeq \Delta_x$.*

Define a partial action $\underline{F}$ of $\underline{M}$ on $\underline{X}$ by $\pi_\bullet(x) \bullet \pi(g) = \pi_\bullet(x \bullet g)$, and, for $\underline{x}$ in $\underline{X}$, let $\Delta_{\underline{x}}$ be the common value of $\pi(\Delta_x)$ for x satisfying $\pi_\bullet(x) = \underline{x}$. Then $\underline{F}$ is a proper partial action of $\underline{M}$ on $\underline{X}$ and $(\Delta_{\underline{x}})_{\underline{x}\in\underline{X}}$ is a right-Garside sequence on $\underline{M}$.

Proof. First, (XI.4.17) and (XI.4.18) guarantee that the definitions of $\underline{x} \bullet \underline{g}$ and $\Delta_{\underline{x}}$ are unambiguous. Next $\underline{F}$ is a partial action of $\underline{M}$ on $\underline{X}$. Indeed, if $\underline{g}$ and $\underline{h}$ lie in $\underline{M}$, and if g, h satisfy $\pi(g) = \underline{g}$ and $\pi(h) = \underline{h}$, then we have $\pi(gh) = \underline{gh}$ and, for $\underline{x}$ in $\underline{X}$ satisfying $\underline{x} = \pi_\bullet(x)$, we can write

$$(\underline{x} \bullet \underline{g}) \bullet \underline{h} = \pi_\bullet(x \bullet g) \bullet \pi(h) = \pi_\bullet((x \bullet g) \bullet h) = \pi_\bullet(x \bullet gh) = \underline{x} \bullet \underline{gh},$$

equality meaning as usual that the involved expressions are simultaneously defined and, in this case, they have the same value.

The partial action $\underline{F}$ is proper. Indeed, assume that $\underline{g}_1, \dots, \underline{g}_m$ are elements of $\underline{M}$. As π is surjective, there exists for every i an element g_i of M that satisfies $\pi(g_i) = \underline{g}_i$. As F is proper, there exists x in X such that $x \bullet g_i$ is defined for each i. Then $\pi_\bullet(x) \bullet \underline{g}_i$ is defined as well.

We now check that $(\Delta_{\underline{x}})_{\underline{x}\in\underline{X}}$ is a right-Garside sequence on $\underline{M}$. First, let $\underline{x}$ belong to $\underline{X}$. There exists x in X satisfying $\pi_\bullet(x) = \underline{x}$. By assumption, $x \bullet \Delta_x$ is defined, hence $\underline{x} \bullet \pi(\Delta_x)$ is defined as well, and it is $\pi_\bullet(x \bullet \Delta_x)$. So the sequence $(\Delta_{\underline{x}})_{\underline{x}\in\underline{X}}$ satisfies (XI.1.11).

Next, the assumption that M is generated by $\bigcup_{x\in X} \mathrm{Div}(\Delta_x)$ implies that $\underline{M}$ is generated by $\bigcup_{x\in X} \pi(\mathrm{Div}(\Delta_x))$, which is $\bigcup_{\underline{x}\in\underline{X}} \mathrm{Div}(\Delta_{\underline{x}})$. So, the sequence $(\Delta_{\underline{x}})_{\underline{x}\in\underline{X}}$ satisfies (XI.1.12).

Now, assume $\underline{s} \in \underline{S}$ and $\underline{s} \preccurlyeq \Delta_{\underline{x}}$. Let s satisfy $\pi(g) = \underline{g}$ and x satisfy $\pi_\bullet(x) = \underline{x}$. By assumption, $\pi_\bullet(x) \bullet \underline{s}$ is defined, hence, by (XI.4.19), $x \bullet \widetilde{\pi}(\underline{s})$ is defined. As $\bigcup_x \mathrm{Div}(\Delta_x)$ generates M, the element $\widetilde{\pi}(\underline{s})$ is left-divisible by some non-invertible element s that lies in $\mathrm{Div}(\Delta_y)$ for some y. As S consists of atoms, every element s of S belongs to $\bigcup_y \mathrm{Div}(\Delta_y)$ and, therefore, by (XI.1.15), $s \preccurlyeq \Delta_x$ must hold for every x such that $x \bullet s$ is defined. Applying this to $\widetilde{\pi}(\underline{s})$, we deduce $\widetilde{\pi}(\underline{s}) \preccurlyeq \Delta_x$, whence, by (XI.1.12), $\Delta_x \preccurlyeq \widetilde{\pi}(\underline{s})\Delta_{x\bullet\widetilde{\pi}(\underline{s})}$. Applying π, we deduce $\Delta_{\underline{x}} \preccurlyeq \underline{s}\Delta_{\underline{x}\bullet\underline{s}}$. Then an easy induction on the length of $\underline{g}$ shows that $\underline{g} \preccurlyeq \Delta_{\underline{x}}$ implies $\Delta_{\underline{x}} \preccurlyeq \underline{g}\Delta_{\underline{x}\bullet\underline{g}}$ for every $\underline{g}$ in $\underline{M}$. So, the sequence $(\Delta_{\underline{x}})_{\underline{x}\in\underline{X}}$ satisfies (XI.1.13).

As of (XI.1.14), it is automatically satisfied by the sequence $(\Delta_{\underline{x}})_{\underline{x}\in\underline{X}}$ since any two elements of $\underline{M}$ are supposed to admit a left-gcd.

Finally, assume that $\underline{g}$ is an element of $\underline{M}$ satisfying $\underline{g} \preccurlyeq \Delta_{\underline{x}}$ and $\underline{y} \bullet \underline{g}$ is defined. Let x, y in X satisfy $\pi_\bullet(x) = \underline{x}$, $\pi_\bullet(y) = \underline{y}$. Let $\underline{w}$ be an $\underline{S}$-word representing $\underline{g}$ in $\underline{M}$. By construction, $[\underline{w}]$ left-divides $\Delta_{\pi_\bullet(x)}$ hence, by (XI.4.20), $[\widetilde{\pi}^*(\underline{w})]$ left-divides Δ_x. By (XI.4.19) and an induction on the length of $\underline{w}$, the assumption that $\underline{y} \bullet [\underline{w}]$ is defined implies that $y \bullet [\widetilde{\pi}^*(\underline{w})]$ is defined. As the sequence $(\Delta_x)_{x\in X}$ satisfies (XI.1.15), we deduce that $[\widetilde{\pi}^*(\underline{w})]$ left-divides Δ_y. Applying π, we conclude that $[\underline{w}]$, that is, $\underline{g}$, left-divides $\Delta_{\underline{y}}$. Hence, the sequence $(\Delta_{\underline{x}})_{\underline{x}\in\underline{X}}$ satisfies (XI.1.15) and, therefore, it is a right-Garside sequence in $\underline{M}$. □

Solution to selected exercises

Exercice 104 (**skeleton**).— *Say that a set of addresses is an* antichain *if it does not contain two addresses, one is a prefix of the other; an antichain is called* maximal *if it is properly included in no antichain.* (i) *Show that a finite maximal antichain is a family $\{\alpha_1, ..., \alpha_n\}$ such that every long enough address admits as a prefix exactly one of the addresses α_i.* (ii) *Show that, for every Σ-word w, there exists a unique finite maximal antichain A_w such that $T \bullet w$ is defined if and only if the skeleton of T includes A_w.*

Solution. (ii) Use induction on the length of w. For w of length one, say $w = \Sigma_\alpha$, the result is true with $T_w^- = T_w^+ = x_1 = \{\alpha 10\}$. Assume now $w = \Sigma_\alpha w'$. It follows from the definition of the action of Σ_α that, for every address γ, there exists a well defined address γ' such that, for every term T such that $T \bullet \Sigma_\alpha$ is defined, the skeleton of $T \bullet \Sigma_\alpha$ contains γ if and only if the skeleton of T contains γ'. Put $\gamma' = \Sigma_\alpha^{-1}(\gamma)$. Then we claim that the result holds with $A_w = \{\alpha 10\} \cup \Sigma_\alpha^{-1}(A_{w'})$. Indeed, $T \bullet w$ is defined if and only if $T \bullet \Sigma_\alpha$ and $(T \bullet \Sigma_\alpha) \bullet w'$ are defined, hence, by induction hypothesis, if and only if the skeleton of T contains $\alpha 10$ and the skeleton of $T \bullet \Sigma_\alpha$ includes $A_{w'}$.

Exercice 105 (**preservation**).— *Assume that M is a left-cancellative monoid and F is a partial action of M on a set X.* (i) *Show that, if M admits right lcms (resp. conditional right-lcms), then so does the category $\mathcal{C}_F(M,X)$.* (ii) *Show that, if M is right-Noetherian, then so is $\mathcal{C}_F(M,X)$.*

Solution. By definition, $(x, f, y) \preccurlyeq (x', f', y')$ in $\mathcal{C}_F(M,X)$ implies $x' = x$ and $f \preccurlyeq f'$ in M. So the assumption that M is right-Noetherian implies that $\mathcal{C}_F(M,X)$ is right-Noetherian as well. Assume that (x, f, y) and (x, g, z) admit a common right-multiple in $\mathcal{C}_F(M,X)$, say $(x, f, y)(y, g', x') = (x, g, z)(z, f', x')$. Then $fg' = gf'$ holds in M. As M is a right-lcm monoid, f and g admit a right-lcm h, and we have $f \preccurlyeq c$, $g \preccurlyeq h$, and $h \preccurlyeq fg'$. By assumption, $x \bullet fg'$ is defined, hence so is $x \bullet h$, and it is obvious to check that $(x, h, x \bullet h)$ is a right-lcm of (x, f, y) and (x, g, z) in $\mathcal{C}_F(M,X)$.

Exercice 107 (**Noetherianity**).— *Assume that F is a proper partial action of a monoid M on a set X and there exists a map $\mu : X \to \mathbb{N}$ such that $\mu(x \bullet g) > \mu(x)$ holds whenever g is not invertible. Show that M is Noetherian and every element of M has a finite height.*

Solution. Let g be a non-invertible element of M, and let $g_1|\cdots|g_p$ be any decomposition of g. At the expense of possibly gathering entries, we may assume that each element g_i is non-invertible. As F is proper, there exists x in X such that

$x \bullet g$ is defined. Then, by (XI.1.3), $x \bullet g_1 \cdots g_i$ is defined for every i. We then obtain $\mu(x) < \mu(x \bullet g_1) < \mu(x \bullet g_1 g_2) < \cdots < \mu(x \bullet g)$, whence $p \leqslant \mu(x \bullet g) - \mu(x)$. So g has a finite height bounded above by $\mu(x \bullet g) - \mu(x)$. In particular, by Proposition II.2.47 (height), M_{LD} is Noetherian.

Exercice 109 (**common multiple**).— *Assume that M is a left-cancellative monoid, F is a partial action of M on X, and $(\Delta_x)_{x \in X}$ is a right-Garside sequence in M with respect to F. Show that, for every x in X, any two elements of* $\mathrm{Def}(x)$ *admit a common right-multiple that lies in* $\mathrm{Def}(x)$.

Solution. Let f, g belong to $\mathrm{Def}(x)$. Then $(x, f, x \bullet f)$ and $(x, g, x \bullet g)$ are two elements of $\mathcal{C}_F(M,X)$ sharing the same source. By Proposition XI.3.25, $\mathcal{C}_F(M,X)$ possesses a right-Garside map, hence any two elements of $\mathcal{C}_F(M,X)$ with the same source admit a common right-multiple. So do in particular $(x, f, x \bullet f)$ and $(x, g, x \bullet g)$. A common right-multiple must be of the form $(x, h, x \bullet h)$ where h is a common right-multiple of f and g in M. Then h lies in $\mathrm{Def}(x)$.

Chapter XII: Ordered groups

Skipped proofs

(none)

Solution to selected exercises

Exercice 111 (**braid ordering**).— *Show that* $1 <_{\mathrm{D}} \sigma_1\sigma_2 \leqslant_{\mathrm{D}} (\sigma_1\sigma_2\sigma_1)^{2p}\sigma_2^q$ *holds in B_3 for $p > 0$.*

Solution. Put $g = (\sigma_1\sigma_2\sigma_1)^{2p}\sigma_2^q$ and $g' = \sigma_1\sigma_2$. We find $g'^{-1}g = \sigma_1(\sigma_1\sigma_2\sigma_1)^{2p-1}\sigma_2^q$. Hence $g' <_{\mathrm{D}} g$ is true.

Exercice 112 (**limit of conjugates**).— *Assuming that $<_{\mathrm{D}}$ is a limit of its conjugates in B_3, show the same result in B_n. [Hint: Use the subgroup of B_n generated by σ_{n-2} and σ_{n-1}, which is isomorphic to B_3.]*

Solution. Let P_n be the positive cone of $<_{\mathrm{D}}$ on B_n, and let H be the subgroup of B_n generated by σ_{n-2} and σ_{n-1}. Then H is isomorphic to B_3 and the $<_{\mathrm{D}}$-ordering of B_n restricted to H corresponds with the $<_{\mathrm{D}}$-ordering of B_3. Moreover, the positive cone for the $<_{\mathrm{D}}$-ordering of H is $P_n \cap H$. Let S be a finite subset of P_n. By assumption, there exists f in H such that $f^{-1}hf$ is lies in P_n for every h in $S \cap H$, and $f(P_n \cap H)f^{-1}$ is distinct from $P_n \cap H$. Assume now $h \in S \setminus H$. By assumption, h is σ_i-positive for some $i < n-2$. Its conjugate $f^{-1}hf$ is σ_i-positive as well, hence it lies in P_n. We deduce $f^{-1}Sf \subseteq P_n$, hence $S \subseteq fP_nf^{-1}$. Finally, fP_nf^{-1} and P_n are distinct, because their intersections with H are distinct.

Exercice 113 (**closure of conjugates**).— *Let P_n be the positive cone of the ordering $<_{\mathrm{D}}$ on B_n considered in Example XII.1.23. Show that the closure of the conjugates of P_n in* $\mathrm{LO}(B_n)$ *is a Cantor set.*

Solution. Let Z_n be the family of all conjugates of P_n. We saw in Example XII.1.23 that P_n is a limit of its conjugates, hence it is a limit point in Z_n. Hence so is every conjugate of P_n, that is, every point of Z_n is a limit point in Z_n. By continuity, every point in the closure of Z_n is a limit point. Next, the closure of Z_n is a closed subspace in a totally disconnected nonempty compact metric space, hence it is itself totally a disconnected, compact, and metric space. As it is nonempty and every point is a limit point, the closure of Z_n is homeomorphic to the Cantor set.

Exercice 114 (**space** $\mathrm{LO}(B_\infty)$).— *Show that every point in the space* $\mathrm{LO}(B_\infty)$ *is a limit of its conjugates and that* $\mathrm{LO}(B_\infty)$ *is homeomorphic to the Cantor set (contrary to the spaces* $\mathrm{LO}(B_n)$ *for finite* n*).*

Solution. Consider an arbitrary positive cone P for a left-invariant ordering of B_∞ and suppose S is a finite subset of P. We will show there is a positive cone $\sigma_i P \sigma_i^{-1}$ in B_∞ that also includes S and is distinct from P. Choose n such that S is included in Bn. Then, for each $i > n$, every braid in S commutes with σ_i, so we have $S = \sigma_i S \sigma_i^{-1} \subseteq \sigma_i P \sigma_i^{-1}$. On the other hand, there exists $i > n$ such that the sets P and $\sigma_i P \sigma_i^{-1}$ are different. For otherwise, using sh for the shift endomorphism that maps σ_i to σ_{i+1} for every i, consider the subgroup $\mathrm{sh}^n(B_\infty)$, which is isomorphic to B_∞. The sets $P \cap \mathrm{sh}^n(B_\infty)$ and $(\sigma_i P \sigma_i^{-1}) \cap \mathrm{sh}^n(B_\infty)$ are positive cones for orderings of $\mathrm{sh}^n(B_\infty)$. If $P = \sigma_i P \sigma_i^{-1}$ is true for each $i > n$, then the cone $P \cap \mathrm{sh}^n(B_\infty)$ of $\mathrm{sh}^n(B_\infty)$ is invariant under conjugation by all elements of $\mathrm{sh}^n(B_\infty)$. This would imply that $\mathrm{sh}^n(B_\infty)$ and, therefore, $B\infty$ are bi-orderable, which is not true.

Exercice 116 (**non-terminating reversing**).— *Assume that* (S, R) *is a triangular presentation.* (i) *Show that, if a relation of* $\widehat{R}$ *has the form* $s = w$ *with* $\lg(w) > 1$ *and* w *finishing with* s*, then the monoid* $\langle S \mid R\rangle^+$ *is not of right-O-type.* (ii) *Let* $(S, \widehat{R})$ *be the maximal right-triangular deduced from* (S, R)*. Show that, if a relation of* $\widehat{R}$ *has the form* $s = w$ *with* w *beginning with* $(uv)^r us$ *with* $r \geqslant 1$*,* u *nonempty, and* v *such that* $v^{-1}s$ *reverses to a word beginning with* s*, hence in particular if* v *is empty or it can be decomposed as* $u_1, \dots, u_m$ *where* $u_k s$ *is a prefix of* w *for every* k*, then* $s^{-1}us$ *cannot be terminating, and deduce that*$\langle S \mid R\rangle^+$ *is not of right-O-type.* (iii) *Show that a relation* $\mathtt{a} = \mathtt{babab}^3\mathtt{a}^2\dots$ *is impossible in a right-triangular presentation for a monoid of right-O-type.*

Solution. (i) If R contains a relation $s = us$ with u nonempty, $s = [u]^+ s$ holds in $\langle S \mid R\rangle^+$, whereas $1 = [u]^+$ fails. So $\langle S \mid R\rangle^+$ is not right-cancellative. (ii) Write the involved relation $s = (uv)^r usw_1$ with $v^{-1}s \curvearrowright_{\widehat{R}} sw_2$. We find

$$\begin{aligned} s^{-1}us \;&\curvearrowright_{\widehat{R}}\; w_1^{-1}s^{-1}(vu)^{-(r-1)}u^{-1}v^{-1}s \\ &\curvearrowright_{\widehat{R}}\; w_1^{-1}s^{-1}(vu)^{-(r-1)}u^{-1}sw_2 \;\curvearrowright_{\widehat{R}} \\ &w_1^{-1}s^{-1}(vu)^{-(r-1)}(vu)^{r-1}vusw_1w_2 \\ &\curvearrowright_{\widehat{R}}\; w_1^{-1}s^{-1}vusw_1w_2 \;\curvearrowright_{\widehat{R}}\; w_1^{-1}w_2^{-1}\cdot s^{-1}us\cdot w_1w_2. \end{aligned}$$

We deduce that $s^{-1}us \;\curvearrowright_{\widehat{R}}\; (w_1^{-1}w_2^{-1})^n \cdot s^{-1}us \cdot (w_1w_2)^n$ holds for every n and, therefore, it is impossible that $s^{-1}us$ leads in finitely many steps to a positive–negative word. Proposition II.4.51 (completeness), right-reversing is complete for

$(S, \widehat{R})$, the elements s and $[u]^+ s$ admit no common right-multiple in $\langle S \mid R\rangle^+$, and the latter cannot be of right-O-type.

(iii) The right-hand side of the relation can be written as $(\mathtt{ba})\mathtt{bab}^2(\mathtt{ba})\mathtt{a}...$, eligible for(ii) with $u = \mathtt{ba}$ and $v = \mathtt{bab} \cdot \mathtt{b}$, a product of two words u_1, u_2 such that $u_i\mathtt{a}$ is a prefix of the right-hand term of the relation.

Exercice 117 (**roots of Garside element**).— *Assume that M is a left-cancellative monoid generated by a set S* (i) *Show that, for δ, g in a left-cancellative monoid M, a necessary and sufficient condition for δ to right-dominate g is that there exist $m \geqslant 1$ satisfying $(*)\forall k \geqslant 0$ ($g\delta^{km+m-1} \preccurlyeq \delta^{km+1}$).* (ii) *Assume that δ^m is a right-Garside element in M. Show that δ right-dominates every element g that satisfies $g\delta^{m-1} \preccurlyeq \delta$.*

Solution. (i) If δ right-dominates g, then, by definition, $(*)$ holds with $m = 1$. Conversely, assume $(*)$. Let n be a nonnegative integer. Let k be maximal with $km \leqslant n$. Then we have $n \leqslant km+m-1$, and $(*)$ implies $g\delta^n \preccurlyeq g\delta^{km+m-1} \preccurlyeq \delta^{km+1} \preccurlyeq \delta^{n+1}$, so δ right-dominates s.

(ii) Put $\Delta = \delta^m$, and let ϕ be the (necessarily unique) endomorphism of M witnessing that Δ is right-quasi-central. First, we have $\delta\Delta = \delta^{m+1} = \Delta\phi(\delta)$, whence $\phi(\delta) = \delta$ since M is left-cancellative. Next, we claim that $g \preccurlyeq h$ implies $\phi(g) \preccurlyeq \phi(h)$. Indeed, by definition, $g \preccurlyeq h$ implies the existence of h' satisfying $gh' = h$, whence $\phi(g)\phi(h') = \phi(h)$ since ϕ is an endomorphism. This shows that $\phi(g) \preccurlyeq \phi(h)$ is satisfied. So, in particular, and owing to the above equality, $g \preccurlyeq \delta$ implies $\phi(g) \preccurlyeq \delta$. Now assume $g\delta^{m-1} \preccurlyeq \delta$. Then, for every k, we find

$$g\delta^{km+m-1} = g\delta^{m-1}\Delta^k = \Delta^k\phi^k(g\delta^{m-1}) \preccurlyeq \Delta^k\phi^k(\delta) = \Delta^k\delta = \delta^{km+1},$$

and we conclude that δ right-dominates g by (i).

Exercice 119 (**right-ceiling**).— *Assume that M is a cancellative monoid of right-O-type, and that $s_\ell \cdots s_1$ is a right-top S-word in M such that $[s_\ell \cdots s_1]^+$ is central in M. Show that $s_i = s_1$ must hold for every i, and deduce that ${}^\infty s_1$ is the right-S-ceiling in M.*

Solution. Let $\Delta = s_\ell \cdots s_1$. First, we have $g \preccurlyeq \Delta$ for every g in S^ℓ, so that Δ right-dominates S^ℓ. Hence, by Lemma XII.3.9, the right-S-ceiling is periodic with period $s_\ell \cdots s_1$. Now consider its length $\ell + 1$ final fragment $s_1 s_\ell \cdots s_1$. Then, in M, we have $s_1 s_\ell \cdots s_1 = s_\ell \cdots s_1 s_1$, so $s_1 s_\ell \cdots s_1$ and $s_\ell \cdots s_1 s_1$ are two right-top S-words of length $\ell + 1$. By uniqueness of the right-S-ceiling, these words must coincide, which is possible only for $s_1 = ... = s_\ell$.

Exercice 123 (**no triangular presentation**]).— *Assume that M is a monoid of right-O-type that is generated by $\mathtt{a}, \mathtt{b}, \mathtt{c}$ with $\mathtt{a} \succ \mathtt{b} \succ \mathtt{c}$ and $\mathtt{b}, \mathtt{c}$ satisfying some relation $\mathtt{b} = \mathtt{c}v$ with no $\mathtt{a}$ in v.* (i) *Prove that, unless M is generated by $\mathtt{b}$ and $\mathtt{c}$, there is no way to complete $\mathtt{b} = \mathtt{c}v$ with a relation $\mathtt{a} = \mathtt{b}u$ so as to obtain a presentation of M.* (ii) *Deduce that no right-triangular presentation made of $\mathtt{b} = \mathtt{cbc}$ (Klein bottle relation) or $\mathtt{b} = \mathtt{cb}^2\mathtt{c}$ (Dubrovina–Dubrovin braid relation) plus a relation of the form $\mathtt{a} = \mathtt{b}...$ may define a monoid of right-O-type.*

Solution. For a contradiction, assume that $(\mathtt{a}, \mathtt{b}, \mathtt{c}; \mathtt{a} = \mathtt{b}u, \mathtt{b} = \mathtt{c}v)$ is a presentation of M. If there is no $\mathtt{a}$ in u, the assumption that $\mathtt{a} = \mathtt{b}u$ is valid in M implies that $\mathtt{a}$ belongs to the submonoid generated by $\mathtt{b}$ and $\mathtt{c}$, so M must be generated

by $\mathtt{b}$ and $\mathtt{c}$.Assume now that there is at least one $\mathtt{a}$ in u. As $\mathtt{a}$ does not occur in $\mathtt{b} = \mathtt{c}v$, a word containing $\mathtt{a}$ cannot be equivalent to a word not containing $\mathtt{a}$. This implies that $\mathtt{a}$ is preponderant in $\{\mathtt{a}, \mathtt{b}, \mathtt{c}\}$. Indeed, assume that g, h belong to the submonoid of M generated by $\mathtt{b}$ and $\mathtt{c}$. By the above remark, $h\mathtt{a}g' = g$ is impossible, hence so is $h\mathtt{a} \preccurlyeq g$. As, by assumption, M is of right-O-type, we deduce $g \preccurlyeq h\mathtt{a}$. Then Proposition XII.3.14 gives the result.

Exercice 124 (**Birman–Ko–Lee generators**).— *Put $b_{i,j} = a_{i,j}^{(-1)^{i+1}}$ in the braid group B_n. Show that, for every n, the monoid $B_n^\oplus$ is generated by the elements $b_{i,j}$.*

Solution. By Lemma XII.3.13, an element of B_n belongs to $B_n^\oplus$ if and only if it is either σ_i-positive for some odd i or σ_i-negative for some even i. The braid relations imply $a_{i,j} = \sigma_{j-1} \cdots \sigma_{i+1} \sigma_i \sigma_{i+1}^{-1} \cdots \sigma_{j-1}^{-1}$ for $i < j$, hence $a_{i,j}$ is σ_i-positive, and $b_{i,j}$ is σ_i-positive for odd i and σ_i-negative for even i. Therefore, $b_{i,j}$ belongs to $B_n^\oplus$ for all i, j. Conversely, in B_n, we have

$$\sigma_i \cdots \sigma_{n-1} = (\sigma_i \cdots \sigma_{n-2} \sigma_{n-1} \sigma_{n-2}^{-1} \cdots \sigma_i^{-1}) \\ (\sigma_i \cdots \sigma_{n-3} \sigma_{n-2} \sigma_{n-3}^{-1} \cdots \sigma_i^{-1}) \cdots (\sigma_i \sigma_{i+1} \sigma_i^{-1})(\sigma_i),$$

whence $s_i = (\sigma_i \cdots \sigma_{n-1})^{(-1)i+1} = b_{i,n} b_{i,n-1} \cdots b_{i,i+1}$ for odd i, and $= b_{i,i+1} \cdots b_{i,n-1} b_{i,n}$ for even i. Hence s_i belongs to the submonoid of B_n generated by the $b_{i,j}$'s and, finally, $B_n^\oplus$ coincides with the latter.

Chapter XIII: Set-theoretic solutions of YBE

Skipped proofs

(none)

Solution to selected exercises

Exercice 125 (**bijective RC-quasigroup**).— *Assume that $(X, \star)$ is an RC-quasigroup. Let $\psi : X \to X$ and $\Psi : X \times X \to X \times X$ be defined by $\psi(a) = a \star a$ and $\Psi(a, b) = (a \star b, b \star a)$. Show that Ψ is injective (resp. bijective) if and only if ψ is.*

Solution. If Ψ is injective, then so is ψ since $\psi(a) = \psi(a')$ implies $\Psi(a, a) = \Psi(a', a')$. On the other hand, if Ψ is surjective, then so is ψ: for every c, there exists (a, b) satisfying $\Psi(a, b) = (c, c)$. As seen in the proof of Lemma XII.2.23(iii), this implies $a = b$, whence $\psi(a) = c$. For the converse, we first compute $(*)$ $\psi(b \star a) = (b \star a) \star (b \star a) = (a \star b) \star (a \star a) = (a \star b) \star \psi(a)$. Assume that ψ is injective and $\Psi(a, b) = \Psi(a', b') = (c, d)$ holds. By $(*)$, we have $c \star \psi(a) = \psi(d) = c \star \psi(a')$. As the left-translation by c and ψ are injective, we deduce $\psi(a) = \psi(a')$, whence $a = a'$, and a symmetric argument gives $b = b'$, so Ψ is injective. Finally, assume that ψ is bijective and c, d belong to X. As the left-translation by c and ψ are surjective, we can find a satisfying $c \star \psi(a) = \psi(d)$, whence, by $(*)$, $\psi(b \star a) = \psi(d)$ and, similarly, we can find b satisfying $\psi(a \star b) = \psi(c)$. As ψ is injective, we deduce $a \star b = c$ and $b \star a = d$, whence $\Psi(a, b) = (c, d)$. So Ψ is surjective, hence bijective.

Exercice 126 (**right-complement**).— *Assume that $(X,\star)$ is an RC-quasigroup and M is the associated structure monoid.* (i) *Show that, for every element f in $M\cap X^p$, the function from X to $X\cup\{1\}$ that maps t to $f\backslash t$ takes pairwise distinct values in X plus at most p times the value 1.* (ii) *Deduce that, for I a finite subset of X with cardinal n, the right-lcm Δ_I of I lies in X^n.*

Solution. (i) We use induction on p. For $p=0$, that is, for $f=1$, we have $f\backslash t=t$, which takes pairwise distinct values in X. For $p=1$, that is, for f in X, we have $f\backslash f=1$ and $f\backslash t=f\star t$ for $t\neq f$. As $t\mapsto f\star f$ injective, the expected result is true. Assume $p\geqslant 2$ and write $f=gs$ with g in X^{p-1}. Then, by the formula for an iterated right-complement, we have $f\backslash t=s\backslash(g\backslash t)$, whence $f\backslash t=1$ if $g\backslash t$ lies in $\{1,s\}$ and $f\backslash t=s\star(g\backslash t)$ otherwise. Then the result follows from the induction hypothesis.

(ii) Use induction on $n\geqslant 0$. For $n=0$, we have $\Delta_I=1$. For $n=1$, say $I=\{s\}$ with s in X, we have $\Delta_I=s$, an element of X. For $n=2$, say $I=\{s,t\}$ with $s\neq t$, we have $\Delta_I=s(s\star t)$, so the induction starts. Assume $n\geqslant 3$ and $I=\{s_1,\dots,s_n\}$. Let $J=\{s_1,\dots,s_{n-2}\}$. By induction hypothesis, Δ_J belongs to X^{n-2}, and $\Delta_{J\cup\{s_{n-1}\}}$, which is $\Delta_J(\Delta_J\backslash s_{n-1})$, belongs to X^{n-1}. This implies $\Delta_J\backslash s_{n-1}\neq 1$, so $\Delta_J\backslash s_{n-1}$ must lie in X. For the same reason, $\Delta_J\backslash s_n$ lies in X. Moreover, as s_{n-1} and s_n are distinct, we have $\Delta_J\backslash s_{n-1}\neq\Delta_J\backslash s_n$ by (i). Now, as I is $J\cup\{s_{n-1}\}\cup\{s_n\}$, Proposition II.2.12 (iterated lcm) gives $\Delta_I=\Delta_{J\cup\{s_{n-1}\}}\cdot(\Delta_J\backslash s_{n-1})\backslash(\Delta_J\backslash s_n)$ with a commutative diagram as below, and the question is to know whether the last term may be 1.



Now we saw above that $\Delta_J\backslash s_{n-1}$ and $\Delta_J\backslash s_n$ are distinct elements of X, so the element $(\Delta_J\backslash s_{n-1})\backslash(\Delta_J\backslash s_n)$ is equal to $(\Delta_J\backslash s_{n-1})\star(\Delta_J\backslash s_n)$, an element of X. Hence Δ_I belongs to X^n, as expected.

Exercice 128 (**I-structure**).— *Assume that $(X,\star)$ is a bijective RC-quasigroup.* (i) *Show (by a direct argument) that the map $\underline{\star}$ from $X^*\times X$ to X defined by $1\underline{\star}t=t$, $s\underline{\star}t=s\star t$ for s in X, and $(u|v)\underline{\star}t=v\underline{\star}(u\underline{\star}t)$ induces a well-defined map from $M\times X$ to X.* (ii) *Show that the map ν from X^* to M defined by $\nu(1)=1$, $\nu(s)=s$, and $\nu(ws)=\nu(w)\cdot(\nu(w)\underline{\star}s)$ for s in X induces a well-defined map from $\mathbb{N}^{(X)}$ to M.*

Solution. (i) Owing to the presentation of M, it suffices to check that the translations associated with $r|(r\star s)$ and $s|(s\star r)$ coincide. Now, using the RC-law, we find $(r|(r\star s))\underline{\star}t=(r\star s)\star(r\star t)=(s\star r)\star(s\star t)=(s|(s\star t))\underline{\star}t$.

(ii) As $\mathbb{N}^{(X)}$ is the quotient of X^* by the equivalence relation generated by all pairs $(u|s|t|v,u|t|s|v)$ with s,t in X, it suffices to check that the images of such words coincide. Now we find

$$\begin{aligned}\nu(u|s|t)&=\nu(u|s)\cdot(\nu(u|s)\underline{\star}t)=\nu(u)\cdot(\nu(u)\underline{\star}s)\cdot(\nu(u)|(\nu(u)\underline{\star}s)\underline{\star}t)\\&=\nu(u)\cdot(\nu(u)\underline{\star}s)\cdot((\nu(u)\underline{\star}s)\underline{\star}(\nu(u)\underline{\star}t))\\&=\nu(u)\cdot(\nu(u)\underline{\star}s)\cdot((\nu(u)\underline{\star}s)\star(\nu(u)\underline{\star}t))\end{aligned}$$

and $\nu(u|t|s)=\nu(u)\cdot(\nu(u)\underline{\star}t)\cdot((\nu(u)\underline{\star}t)\star(\nu(u)\underline{\star}s))$, whence the equality. Then multiplying by v on the right gives the same result.

Exercice 129 (**parabolic submonoid**).— (i) *Assume that (X, ρ) is a finite involutive nondegenerate set-theoretic solution of YBE and M is the associated structure monoid. Show that a submonoid M_1 of M is parabolic if and only if there exists a (unique) subset I of X satisfying $\rho(I \times I) = I \times I$ such that M_1 is the submonoid of M generated by I.* (ii) *Assume that $(X, \star)$ is a finite RC-system and M is the associated structure monoid. Then a submonoid M_1 of M is parabolic if and only if there exists a (unique) subset I of X such that I is closed under $\star$ and M_1 is the submonoid generated by I.* (iii) *Show that, if $(X, \star)$ is an infinite RC-quasigroup, there may exist subsets I of X that are closed under $\star$ but the induced RC-system $(I, \star)$ is not an RC-quasigroup.*

Solution. (i) Assume that M_1 is a parabolic submonoid of M. Set $I = M_1 \cap X$. Since X generates M and M_1 is closed under factors, M_1 is generated by I. Now, assume that a, b belong to I and $(c, d) = \rho(a, b)$ holds. Then ab belongs to M_1, and $ab = cd$ holds in M. As M_1 is closed under factor, we deduce that c and d lie in I, whence $\rho(I \times I) \subseteq I \times I$. As ρ is involutive, the latter inclusion must be an equality. Conversely assume I is a subset of X satisfying $\rho(I \times I) = I \times I$ and M_1 is the submonoid of M generated by I. The monoid is (right)-Noetherian, so, in order to prove that M_1 is a parabolic submonoid, and owing to Proposition VII.1.32 (parabolic subcategory), it is sufficient to establish that M_1 is closed under factor and right-comultiple, that is, in a context where right-lcms exist, that M_1 is closed under factor and right-lcm. Now, an obvious induction shows that every X-word that is equivalent to an I-word is itself an I-word, implying that M_1 is closed under factor. On the other hand, let ρ_I be the restriction of ρ to $I \times I$. The assumption that (X, ρ) is a set-theoretic solution of YBE implies that so is (I, ρ_I), and the assumption that (X, ρ) is involutive implies that so is (I, ρ_I). Finally, the assumption that (X, ρ) is nondegenerate implies that so is (I, ρ_I): with the notation of Definition XIII.1.8, the left-translations associated with ρ_1 and right-translations associated with ρ_2 are injective on X, hence so are their restrictions to I, and therefore, the latter are bijective since I is finite. Now, let a, b belong to I. As (I, ρ_I) is an involutive nondegenerate solution of YBE, there exist c, d in I satisfying $ac = bd$ in M, hence in M_1, and M_1 is closed under right-lcm in M. Hence M_1 is a parabolic submonoid of M.

(ii) Translate the result of (i) in terms of the operation $\star$.

(iii) Consider $X = \mathbb{Z}$ and $x \star y = y + 1$. Then the restriction of $\star$ to $\mathbb{N}$ does not give an RC-quasigroup.

Chapter XIV: More examples

Skipped proofs

(none)

Solution to selected exercises

(none)

Laboratoire de Mathématiques Nicolas Oresme, CNRS UMR 6139, Université de Caen, 14032 Caen, France
E-mail address: patrick.dehornoy@unicaen.fr
URL: www.math.unicaen.fr/∼dehornoy

Laboratoire Amiénois de Mathématique Fondamentale et Appliquée, CNRS UMR 7352, Université de Picardie Jules-Verne, 80039 Amiens, France
E-mail address: digne@u-picardie.fr
URL: www.mathinfo.u-picardie.fr/digne/

Laboratoire de Mathématiques Nicolas Oresme, CNRS UMR 6139, Université de Caen, 14032 Caen, France
E-mail address: godelle@math.unicaen.fr
URL: www.math.unicaen.fr/∼godelle

Mathematics Institute, University of Warwick, Coventry CV4 7AL, United Kingdom
E-mail address: D.Krammer@warwick.ac.uk
URL: www.warwick.ac.uk/∼masbal/

Institut de Mathématiques de Jussieu, CNRS UMR 7586, Université Denis Diderot Paris 7, 75205 Paris 13, France
E-mail address: jmichel@math.jussieu.fr
URL: www.math.jussieu.fr/∼jmichel/